

The near-flat-space and BMN limits for strings in $AdS_4 \times CP^3$ at one loop

Michael C Abbott¹ and Per Sundin²

¹ Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Rd, Mumbai 400 005, India

² Astrophysics, Cosmology & Gravity Centre and Department of Applied Mathematics, University of Cape Town, Private Bag, Rondebosch, 7700, South Africa

E-mail: abbott@theory.tifr.res.in and nidnus.rep@gmail.com

Received 27 August 2011, in final form 8 November 2011

Published 8 December 2011

Online at stacks.iop.org/JPhysA/45/025401

Abstract

This paper studies type IIA string theory in $AdS_4 \times CP^3$ in both the Berenstein–Maldacena–Nastase (BMN) and Maldacena–Swanson or near-flat-space limits. We derive the simpler Lagrangian for the latter limit by taking a large worldsheet boost of the BMN theory. We then calculate one-loop corrections to the correlators of the various fields using both theories. In all cases the near-flat-space results agree with a limit of the BMN results, providing evidence for the quantum consistency of this truncation. The corrections can also be compared to an expansion of the exact dispersion relation, known from integrability apart from one interpolating function $h(\lambda)$. Here we see agreement with the results of McLoughlin *et al* (2008 *J. High Energy Phys.* JHEP11(2008)069), and observe that it does not appear to be possible to fully implement the cutoff suggested by Gromov & Mikhaylov (2009 *J. High Energy Phys.* JHEP04(2009)083), although for some terms we can do so. In both the near-flat-space and BMN calculations there are some extra terms in the mass shifts which break supersymmetry. These terms are extremely sensitive to the cutoff used, and can perhaps be seen as a consequence of using dimensional regularization.

PACS numbers: 11.25.Tq, 11.25.Db, 11.15.Yc

1. Introduction

Closed strings in flat spacetime have the property that left- and right-moving excitations decouple, but this is no longer true in curved backgrounds such as $AdS_5 \times S^5$ or $AdS_4 \times CP^3$. Even after one has taken the Berenstein–Maldacena–Nastase (BMN), or Penrose limit, focusing on strings very near to a light-like trajectory [5], nontrivial couplings between the left movers and right movers remain. The limit proposed by Maldacena and Swanson in [1] can

be viewed as a partial restoration of this decoupling, by virtue of the fact that one light-like worldsheet momentum is taken to be much larger than the other, $p_- \gg p_+$.

Both of these limits are taken using a power of the 't Hooft coupling λ as the parameter, thus tying them to the semiclassical limit $\lambda \gg 1$. The BMN limit looks at perturbations of order $1/\lambda^{1/4}$ from a null geodesic, allowing us (at a given loop order) to truncate the expansion of the Lagrangian in the number of fields. The near-flat-space limit then scales the momenta such that $p_-/p_+ \sim \sqrt{\lambda}$. This can be regarded as a worldsheet boost in the σ^- direction, and the simplification comes from discarding all interaction terms except the leading ones under this boost. It is not obvious that this truncation will be respected quantum mechanically. For this to happen, the contributions from where momenta on internal lines are *not* large must cancel out, or at least be sufficiently small.

In the case of strings in $AdS_5 \times S^5$, this quantum consistency was checked at one loop in [6] and at two loops in [7]. These papers computed primarily corrections to four-point functions and compared these to the worldsheet S -matrix known exactly from integrability [8]. The second paper [7] also computed corrections to the two-point function and compared these to the dispersion relation $E(p)$. It is useful here to think of the near-flat-space limit as being intermediate between the BMN limit and the full sigma model:

$$\begin{aligned} p &\sim 1/\sqrt{\lambda}, & \text{BMN limit} \\ p &\sim \lambda^{-1/4}, & \text{near-flat-space} \\ p &\sim 1, & \text{giant magnons.} \end{aligned} \tag{1}$$

We write ‘giant magnons’ for the third sector as these are the classical string solutions obeying the dispersion relation at finite p [9]. They are of size $\Delta\phi = p$ along an equator of S^5 .

For the case of strings in $AdS_4 \times CP^3$, relevant for the comparison with ABJM’s superconformal Chern–Simons theory [10, 11], only the bosonic part of the near-flat-space limit has been studied [12]. Fermions are of course essential for one-loop calculations, and so we work out the complete near-flat-space Lagrangian, starting from the BMN case of [13]. We use this to calculate corrections to two-point functions, which we can compare to the exact dispersion relation

$$E = \sqrt{\frac{1}{4} + 4h(\lambda)^2 \sin^2 \frac{p}{2}}. \tag{2}$$


The interpolating function $h(\lambda)$ is the only part of this relation not fixed by symmetries. At small values of λ it is $h(\lambda) = \lambda + \mathcal{O}(\lambda^3)$, but at large λ (relevant for semiclassical strings) it is instead

$$h(\lambda) = \sqrt{\frac{\lambda}{2}} + c + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right). \tag{3}$$

The subleading term c has been the subject of some debate in the literature. In one-loop energy corrections to spinning strings [14–16, 3] and to giant magnons [17], there is a choice of how to regulate divergent sums over mode numbers, and using the same momentum cutoff Λ for all modes leads to $c = -\log 2/2\pi$. However, it has been argued that it is natural to use a cutoff of 2Λ for half of the modes (the heavy modes) and this ‘new’ prescription leads instead to $c = 0$ [3, 18]. The same two prescriptions have been implemented in a recent Hamiltonian analysis [4], which argues in favour of the new prescription. Our paper is in some ways the Lagrangian analysis complementary to this.

The sigma model in $AdS_4 \times CP^3$ has several novel features compared to that in $AdS_5 \times S^5$. One is that its excitations do not all have the same mass: half are ‘light’ ($m = \frac{1}{2}$ in our conventions) and half are ‘heavy’ ($m = 1$). Another is that the Lagrangian has interactions starting at cubic order, rather than at quartic order [19, 12, 20], greatly expanding the number


of Feynman diagrams possible. The cubic interactions always couple two light modes to one heavy mode, and therefore lead to bubble diagrams such as



where we draw the heavy mode as a double line. In this diagram, the same loop momentum applies to both the heavy and the light modes, at least in the UV. This points towards the ‘old’ prescription of integrating up to the same momentum Λ for all modes. In this paper, we use primarily dimensional regularization, but for the simplest case can also obtain exactly the same results using a momentum cutoff.

In addition to the term coming from $c = -\log 2/2\pi$, we also find (using either regularization procedure) some additional terms. These terms are present in both the BMN and near-flat-space limits, and so do not represent a problem with the consistency of the near-flat-space truncation. These extra terms differ for the various particles, vanishing in the case of corrections to the light fermion propagator. This implies that they break supersymmetry, possibly as a consequence of dimensional regularization.

In fact, these extra terms come (primarily) from the bubble diagram above, while the term containing $c = -\log 2/2\pi$ comes from the tadpole diagrams:



Each loop here contains only one mode, and so for these tadpole diagrams there is no obstruction to changing the heavy-mode cutoff to implement the ‘new’ sum. This change cancels the $-\log 2/2\pi$ term, and thus while slightly unsatisfying (since it does nothing to the bubble diagrams) could be said to lead to $c = 0$.

In this paper, we also further the investigation of what one-loop corrections can teach us about the nature of the heavy modes, as initiated by Zarembo in [21] and continued in [13]. There it was argued that the heavy modes dissolve into a multi-particle continuum, under the assumption that $c = 0$. In our calculation, it is clear that the decay of a heavy mode into two light modes is kinematically allowed.

Outline

In section 2, we set up the theory we consider, re-write some results of [13] for the BMN limit in more convenient notation and take the near-flat space limit of the Lagrangian.

Sections 3–5 study mass corrections to the propagator in the near-flat space limit. The basic light boson calculation is quite simple, but the light fermion case is more involved. We discuss some issues about momentum cutoffs in section 4 before working on the heavy modes in section 5.

We then turn to the full near-BMN case in section 6. All of the same issues arise here, but the calculations are a great deal more complicated and thus less transparent. We can reproduce our near-flat-space results as limits of these results. We conclude and summarize in section 7.

Appendix A contains extra notation particularly about fermions. Appendix B contains manipulations to simplify \mathcal{L}_4 . Appendix C looks at expansions of integrals used for dimensional regularization.

2. The string Lagrangian and two strong-coupling limits

Our starting point is the gauge-fixed Lagrangian for type IIA strings in the near-BMN limit of $AdS_4 \times CP^3$ as derived in [13]. We begin by reviewing this derivation very briefly.

We are interested in strings moving in the quotient super-space [22, 23]

$$\frac{OSP(2, 2|6)}{SO(3, 1) \times U(3)},$$

the bosonic part of which is $AdS_4 \times CP^3$. The sigma model is defined in terms of the following Lie-algebra-valued flat current:

$$\begin{aligned} A_\mu(\sigma, \tau) &= -G^{-1} \partial_\mu G, \quad G \in OSP(2, 2|6) \\ &= A_\mu^{(0)} + A_\mu^{(1)} + A_\mu^{(2)} + A_\mu^{(3)}. \end{aligned}$$

Here, $A^{(k)}$ is the component with eigenvalue i^k under the generator of the \mathbb{Z}_4 automorphism, Ω . The subalgebra $so(3, 1) \oplus u(3)$ is precisely that fixed by Ω , and thus to be omitted from the action. The $k = 0, 2$ components are bosonic, and $k = 1, 3$ components are fermionic. The action defined by [22] is as follows:

$$S = \frac{g}{2} \int d^2\sigma \text{Str}[\sqrt{-h} h^{\mu\nu} A_\mu^{(2)} A_\nu^{(2)} + \kappa \epsilon^{\mu\nu} A_\mu^{(1)} A_\nu^{(3)}]. \quad (4)$$

The coupling constant is $g = \sqrt{\lambda/2} = R^2/8\pi\alpha'$, which is taken to be large. In order for local κ -symmetry to hold we need $\kappa^2 = 1$, and we now choose $\kappa = 1$.³

It is necessary to introduce some parameterization of the group. Since we are interested in strings near to the null line $\phi = t$, and will later discard these two directions during gauge fixing, it has been found to be convenient to factorize them out from the start. A suitable parameterization is [13]

$$G = \Lambda(t, \phi) F(\chi) G_\perp(X).$$

The three factors contain the light-cone directions, the fermions and the eight transverse bosonic directions. Explicitly,

$$\begin{aligned} \Lambda(t, \phi) &= \exp \left[\frac{i}{2} \left(x^+ + \left(\frac{1}{2} - a \right) x^- \right) \Sigma_+ + \frac{i}{4} x^- \Sigma_- \right] \\ G_\perp(X) &= G_{AdS} \oplus G_{CP} = \frac{1 + \frac{i}{2} z_i \Gamma^i}{\sqrt{1 - z_i^2/4}} \oplus \exp \left(W + \bar{W} + \frac{1}{2} y T_5 \right) \end{aligned} \quad (5)$$

$$F(\chi) = \chi + \sqrt{1 + \chi^2}.$$

Here, z_i and t are the AdS_4 coordinates, while y , W , \bar{W} and ϕ are the CP^3 coordinates. All the fermionic fields are contained in the matrix χ ; see appendix A for details. The target-space light-cone coordinates here are non-standard ones, defined as

$$x^+ = (1 - a)t + a\phi, \quad x^- = \phi - t,$$

where $a \in [0, 1]$ is the same constant as appears in the generalized light-cone gauge.

The bosonic gauge fixing is simplified by introducing an auxiliary field Π conjugate to $A^{(2)}$. This brings us to a first-order formalism in which the Weyl-invariant worldsheet metric $\gamma^{\mu\nu} = \sqrt{-\det h} h^{\mu\nu}$ enters as a set of Lagrange multipliers:

$$S = -g \int d\sigma \text{Str} \left[\Pi A_0 + \frac{1}{2} \epsilon^{\alpha\beta} A_\alpha^{(1)} A_\beta^{(3)} - \frac{1}{2\gamma^{00}} (\Pi^2 + (A_1^{(2)})^2) + \frac{\gamma^{01}}{\gamma^{00}} \Pi A_1^{(2)} \right]. \quad (6)$$

By solving for Π this action is classically equivalent to (4); for details see [13]. The gauge we adopt is the generalized light-cone gauge of [24, 25], in which we set $x^+ = \tau$ and the density of the momentum p_+ to be a constant:

$$P_+ = g \int d\sigma p_+ = (1 - a)J + a\Delta,$$

³ κ changes the sign under $\sigma \rightarrow -\sigma$.

where Δ and J are the conserved charges from AdS_4 and CP^3 . Note that this gauge relates the length of the string worldsheet to P_+ and the coupling. As a result of this, the large coupling expansion will decompactify the worldsheet. The light-cone gauge also implies that the physical Hamiltonian is

$$\mathcal{H}_{lc} = -P_- = g \int d\sigma p_- = J - \Delta.$$

The gauge fixing breaks some of the symmetries and the set of charges that commutes with the light-cone Hamiltonian combines into $SU(2|2) \times U(1)$ [26].

What we have written thus far is exact in the sense that arbitrarily large motions of the string are allowed. We now take the BMN, or PP-wave limit, in which we focus on strings near to the null geodesic $\phi = t$. By ‘near’ we mean of order $1/\sqrt{g}$, which means that this limit is to be taken simultaneously with the semiclassical limit. We implement this by scaling all fields, making the following replacements in \mathcal{L} :

$$x \rightarrow \frac{1}{\sqrt{g}}x, \quad \pi \rightarrow \frac{1}{\sqrt{g}}\pi, \quad \chi \rightarrow \frac{1}{\sqrt{g}}\chi, \quad (7)$$

writing x for a generic boson (whose conjugate momentum is π) and χ for a fermion. At large g , the Lagrangian is then organized as an expansion in the number of fields, which we write as follows:

$$S = \int d^2\sigma \left[\mathcal{L}_2(\pi, x, \chi) + \frac{1}{\sqrt{g}}\mathcal{L}_3(\pi, x, \chi) + \frac{1}{g}\mathcal{L}_4(\pi, x, \chi) + \mathcal{O}(g^{-3/2}) \right].$$

The result we take from [13] as our starting point is the Lagrangian in this form. (The only change is that we generalize it to the gauge $a \neq \frac{1}{2}$.) Note that unlike the Hamiltonian analysis of [13, 27], it is not necessary for us to perform complicated redefinitions of the fermions in order to make canonically conjugate pairs (up to some order). We can simply work with them as they stand.

2.1. The BMN Lagrangian

The analysis we want to perform in this paper is simplified by eliminating the momentum variables (introduced for the ease of gauge fixing) in favour of velocities. The equations of motion for the momenta π are, schematically,

$$\frac{\partial \mathcal{L}}{\partial \pi} = 0 \quad \Rightarrow \quad \pi = \partial_0 x + \frac{1}{\sqrt{g}} \frac{\partial \mathcal{L}_3}{\partial \pi} + \frac{1}{g} \frac{\partial \mathcal{L}_4}{\partial \pi} + \dots$$

The higher order terms on the right mean that the interaction terms \mathcal{L}_3 and \mathcal{L}_4 will change in nontrivial ways. But the quadratic Lagrangian is essentially⁴ given by replacing $\pi \rightarrow \partial_0 x$ in that of [13]

$$\begin{aligned} \mathcal{L}_2 = & \frac{1}{2} \partial_+ y \partial_- y + \frac{1}{2} \partial_+ z_i \partial_- z_i + \frac{1}{4} \partial_+ \omega_\alpha \partial_- \bar{\omega}^\alpha + \frac{1}{4} \partial_- \omega_\alpha \partial_+ \bar{\omega}^\alpha - \frac{1}{2} (y^2 + z_i^2) - \frac{1}{8} \omega_\alpha \bar{\omega}^\alpha \\ & + i (\bar{\psi}_{+a} \partial_- \psi_+^a + \bar{\psi}_{-a} \partial_+ \psi_-^a) + \frac{i}{2} [(s_-)_\alpha^a \partial_+ (s_-)_a^\alpha + (s_+)_\alpha^a \partial_- (s_+)_a^\alpha] \\ & - \frac{1}{2} (\bar{\psi}_{-a} \psi_+^a + \bar{\psi}_{+a} \psi_-^a) - i (s_+)_\alpha^a (s_-)_a^\alpha. \end{aligned} \quad (8)$$

Here we have introduced worldsheet light-cone coordinates

$$\sigma^\pm = \frac{\tau \pm \sigma}{2}, \quad \Rightarrow \quad \partial_\pm = \partial_0 \pm \partial_1.$$

⁴ We detail some changes of notation in appendix A.

We will refer to both fields and derivatives with a + subscript as being left moving, and those with − as right moving. The list of fields is as follows:

$$\begin{array}{lll}
 \psi_{\pm}^a & \text{complex light fermion} & a = 1, 2 \\
 (s_{\pm})_{\alpha}^a & \text{real heavy fermion} & \alpha = 3, 4 \\
 \omega_{\alpha} & \text{complex light boson} & \\
 y, z_i & \text{real heavy boson} & i = 1, 2, 3.
 \end{array} \tag{9}$$

The fields transform covariantly under the bosonic part of $SU(2|2) \times U(1)$, namely $SU(2)^2 \times U(1)$, and we denote the $SU(2)$ from AdS_4 with Latin indices and the $SU(2)$ from CP_3 with Greek. Both are raised and lowered with ϵ -tensors:

$$R^a = \epsilon^{ab} R_b = \epsilon^{ab} (\epsilon_{bc} R^c), \quad \epsilon_{12} = \epsilon_{34} = 1 \quad \Rightarrow \quad \epsilon^{12} = \epsilon^{34} = -1$$

and the action of conjugation is $(C_{\alpha}^{\beta})^{\dagger} = \bar{C}_{\beta}^{\alpha}$. The light fields also transform under the $U(1)$, with ψ_{\pm} and ω_{α} having +1 charge.

Either by direct inspection or through solving the quadratic equations of motion, one can derive the following Feynman propagators:

$$\langle \bar{\omega}^{\alpha} \omega_{\beta} \rangle = \delta_{\beta}^{\alpha} \frac{2i}{p^2 - \frac{1}{4}}, \quad \langle yy \rangle = \frac{i}{p^2 - 1}, \quad \langle z_i z_j \rangle = \delta_{ij} \frac{i}{p^2 - 1}. \tag{10}$$

(Note that our light boson has non-standard normalization.) For the fermions,

$$\begin{aligned}
 \langle \bar{\psi}_a \psi^b \rangle &= \delta_a^b \frac{D_{\psi\psi}}{p^2 - \frac{1}{4}}, \quad \text{where } D_{\psi\psi} = i p_{+} \text{ for the case } \bar{\psi}_{+} \psi_{+} \\
 &= i p_{-} \quad \quad \quad -, - \\
 &= -1 \quad \quad \quad +, - \\
 &= +1 \quad \quad \quad -, +
 \end{aligned} \tag{11}$$

and

$$\begin{aligned}
 \langle s_{\alpha}^{\beta} s_b^{\beta} \rangle &= \delta_b^{\alpha} \delta_{\alpha}^{\beta} \frac{D_{ss}}{p^2 - 1}, \quad D_{ss} = i p_{+} \text{ for } s_{+} s_{+} \\
 &= i p_{-} \quad \quad \quad -, - \\
 &= -i/2 \quad \quad \quad +, - \text{ or } -, +.
 \end{aligned}$$

The cubic Lagrangian is naturally more complicated, and writing $\overleftrightarrow{\partial}_{\pm} = \overleftarrow{\partial}_{\pm} - \overrightarrow{\partial}_{\pm}$ and $Z_b^a = \sum_i z_i (\sigma_i)_{\beta}^a$, it is given by

$$\begin{aligned}
 \mathcal{L}_3 &= \frac{i}{8} y (\omega_{\alpha} \overleftrightarrow{\partial}_{+} \bar{\omega}^{\alpha} + \omega_{\alpha} \overleftrightarrow{\partial}_{-} \bar{\omega}^{\alpha}) + \frac{i}{2} \bar{\psi}_{-a} \overleftrightarrow{\partial}_{+} \psi_{-}^b \partial_{-} Z_b^a - \frac{i}{2} \bar{\psi}_{+a} \overleftrightarrow{\partial}_{-} \psi_{+}^b \partial_{+} Z_b^a \\
 &+ \frac{1}{2} (\bar{\psi}_{-a} \partial_{+} \psi_{+}^b + \partial_{+} \bar{\psi}_{+a} \psi_{-}^b - \bar{\psi}_{+a} \partial_{-} \psi_{-}^b - \partial_{-} \bar{\psi}_{-a} \psi_{+}^b) Z_b^a \\
 &- \epsilon^{ab} \left[\left(\frac{3i}{16} (s_{-})_{\alpha}^a \bar{\psi}_{-b} + \frac{3}{16} (s_{+})_{\alpha}^a \bar{\psi}_{+b} + \frac{i}{2} (s_{+})_{\alpha}^a \partial_{-} \bar{\psi}_{-b} \right. \right. \\
 &- \frac{1}{2} (s_{-})_{\alpha}^a \partial_{+} \bar{\psi}_{+b} - \frac{1}{4} \partial_{-} (s_{-})_{\alpha}^a \bar{\psi}_{+b} - \frac{i}{4} \partial_{-} (s_{-})_{\alpha}^a \partial_{+} \bar{\psi}_{-b} + \frac{i}{4} \partial_{+} (s_{-})_{\alpha}^a \partial_{-} \bar{\psi}_{-b} \\
 &+ \frac{i}{4} \partial_{+} (s_{+})_{\alpha}^a \bar{\psi}_{-b} + \frac{1}{4} \partial_{-} (s_{+})_{\alpha}^a \partial_{+} \bar{\psi}_{+b} - \frac{1}{4} \partial_{+} (s_{+})_{\alpha}^a \partial_{-} \bar{\psi}_{+b} \Big) \omega_{\alpha} \\
 &- \left(\frac{i}{8} (s_{-})_{\alpha}^a \overleftrightarrow{\partial}_{-} \bar{\psi}_{-b} + \frac{1}{8} (s_{+})_{\alpha}^a \overleftrightarrow{\partial}_{-} \bar{\psi}_{+b} \right) \partial_{+} \omega_{\alpha} \\
 &- \left. \left(\frac{i}{8} (s_{-})_{\alpha}^a \overleftrightarrow{\partial}_{+} \bar{\psi}_{-b} + \frac{1}{8} (s_{+})_{\alpha}^a \overleftrightarrow{\partial}_{+} \bar{\psi}_{+b} \right) \partial_{-} \omega_{\alpha} \right]
 \end{aligned}$$

$$\begin{aligned}
& + \epsilon_{ab} \left[\frac{3i}{16} (s_-)_\alpha^a \psi_-^b - \frac{3}{16} (s_+)_\alpha^a \psi_+^b + \frac{i}{2} (s_+)_\alpha^a \partial_- \psi_-^b \right. \\
& + \frac{1}{2} (s_-)_\alpha^a \partial_+ \psi_+^b + \frac{1}{4} \partial_- (s_-)_\alpha^a \psi_+^b - \frac{i}{4} \partial_- (s_-)_\alpha^a \partial_+ \psi_-^b + \frac{i}{4} \partial_+ (s_-)_\alpha^a \partial_- \psi_-^b \\
& + \frac{i}{4} \partial_+ (s_+)_\alpha^a \psi_-^b - \frac{1}{4} \partial_- (s_+)_\alpha^a \partial_+ \psi_+^b + \frac{1}{4} \partial_+ (s_+)_\alpha^a \partial_- \psi_+^b \left. \right) \bar{\omega}^\alpha \\
& - \left(\frac{i}{8} (s_-)_\alpha^a \overset{\leftrightarrow}{\partial} - \psi_-^b - \frac{1}{8} (s_+)_\alpha^a \overset{\leftrightarrow}{\partial} - \psi_+^b \right) \partial_+ \bar{\omega}^\alpha \\
& - \left(\frac{i}{8} (s_-)_\alpha^a \overset{\leftrightarrow}{\partial} + \psi_-^b - \frac{1}{8} (s_+)_\alpha^a \overset{\leftrightarrow}{\partial} + \psi_+^b \right) \partial_- \bar{\omega}^\alpha \left. \right]. \tag{12}
\end{aligned}$$

The quartic BMN Lagrangian is very involved and we will not present it here.

2.2. The near-flat-space limit

In the near-flat-space limit we focus on those terms which are important at large p_- . We can do this by taking a large worldsheet boost (in the σ^- direction) and keeping only the leading terms under this boost. The quadratic Lagrangian is Lorentz invariant and so is unaffected, but the cubic and quartic interaction terms break this symmetry, and are thus simplified in this limit. These simplifications are the main reason for studying this limit.

The bosonic fields all behave trivially under worldsheet Lorentz transformations. We have written the fermionic fields in terms of left- and right-moving components, and it is easy to see that these must scale like $\sqrt{p_\pm}$ for \mathcal{L}_2 to be invariant. Explicitly, the boost involves the following replacements in \mathcal{L} :

$$\begin{aligned}
\partial_\pm & \rightarrow g^{\mp 1/2} \partial_\pm \quad (\text{i.e. } \sigma^\pm \rightarrow g^{\pm 1/2} \sigma^\pm) \\
\psi_\pm & \rightarrow g^{\mp 1/4} \psi_\pm \quad \text{and likewise } s_\pm \rightarrow g^{\mp 1/4} s_\pm.
\end{aligned} \tag{13}$$

The leading behaviour of \mathcal{L}_3 is given by terms that grow like \sqrt{g} , and we keep only these terms. Likewise the leading terms in \mathcal{L}_4 grow like g . We write the result of these replacements as follows:

$$\frac{1}{\sqrt{g}} \mathcal{L}_3 + \frac{1}{g} \mathcal{L}_4 \rightarrow \mathcal{L}_3^{\text{NFS}} + \mathcal{L}_4^{\text{NFS}} + \mathcal{O}\left(\frac{1}{\sqrt{g}}\right),$$

and after discarding the $\mathcal{O}(1/\sqrt{g})$ terms are thus studying

$$S = \int d^2\sigma [\mathcal{L}_2 + \mathcal{L}_3^{\text{NFS}} + \mathcal{L}_4^{\text{NFS}}]. \tag{14}$$

As noted by [1], this action has no parameters at all.

The idea of [6] is to perform a second boost of the same form:

$$\partial_\pm \rightarrow \gamma^{\mp 1/2} \partial_\pm, \quad \psi_\pm \rightarrow \gamma^{\mp 1/4} \psi_\pm, \quad s_\pm \rightarrow \gamma^{\mp 1/4} s_\pm, \tag{15}$$

leading to

$$S = \int d^2\sigma [\mathcal{L}_2 + \sqrt{\gamma} \mathcal{L}_3^{\text{NFS}} + \gamma \mathcal{L}_4^{\text{NFS}}]. \tag{16}$$

If γ is kept arbitrary, it can be viewed as just a parameter to keep track of the orders. But if $\gamma = 1/g$ (as we will assume), then this second transformation is the inverse of the first, except for the fact that we do not recover the interaction terms we discarded. So what has happened is that we are back in the original variables, but have built in the assumption that $p_- \gg p_+$.

We now write the interaction terms, starting with the cubic term. Here we simply keep those terms in (12) which grow as \sqrt{g} under the boost (13) (i.e. grow as fast as the right-moving momentum p_-):

$$\begin{aligned} \mathcal{L}_3^{\text{NFS}} = & \frac{i}{8} y \omega_\alpha \overleftrightarrow{\partial}_- \bar{\omega}^\alpha + \frac{i}{2} \bar{\psi}_{-a} \overleftrightarrow{\partial}_+ \psi_-^b \partial_- Z_b^a - \frac{1}{2} (\bar{\psi}_{+a} \partial_- \psi_-^b + \partial_- \bar{\psi}_{-a} \psi_+^b) Z_b^a \\ & - \epsilon^{ab} \left[\left(\frac{3i}{16} (s_-)_a^\alpha \bar{\psi}_{-b} + \frac{i}{2} (s_+)_a^\alpha \partial_- \bar{\psi}_{-b} - \frac{1}{4} \partial_- (s_-)_a^\alpha \bar{\psi}_{+b} \right. \right. \\ & \left. \left. - \frac{i}{4} \partial_- (s_-)_a^\alpha \partial_+ \bar{\psi}_{-b} + \frac{i}{4} \partial_+ (s_-)_a^\alpha \partial_- \bar{\psi}_{-b} \right) \omega_\alpha \right. \\ & \left. - \frac{i}{8} (s_-)_a^\alpha \overleftrightarrow{\partial}_- \bar{\psi}_{-b} \partial_+ \omega_\alpha - \frac{i}{8} (s_-)_a^\alpha \overleftrightarrow{\partial}_+ \bar{\psi}_{-b} \partial_- \omega_\alpha \right] \\ & + \epsilon_{ab} \left[\left(\frac{3i}{16} (s_-)_a^\alpha \psi_-^b + \frac{i}{2} (s_+)_a^\alpha \partial_- \psi_-^b + \frac{1}{4} \partial_- (s_-)_a^\alpha \psi_+^b - \frac{i}{4} \partial_- (s_-)_a^\alpha \partial_+ \psi_-^b \right. \right. \\ & \left. \left. + \frac{i}{4} \partial_+ (s_-)_a^\alpha \partial_- \psi_-^b \right) \bar{\omega}^\alpha - \frac{i}{8} (s_-)_a^\alpha \overleftrightarrow{\partial}_- \psi_-^b \partial_+ \bar{\omega}^\alpha - \frac{i}{8} (s_-)_a^\alpha \overleftrightarrow{\partial}_+ \psi_-^b \partial_- \bar{\omega}^\alpha \right]. \end{aligned} \quad (17)$$

We note immediately a distinction from the $AdS_5 \times S^5$ case; this leading-order interaction term has both left- and right-moving fields and derivatives, rather than consisting only of right-moving objects (ψ_- , s_- and ∂_-) as in [1]. This is not unexpected from the form of the boost (13); in order for a term with two right-moving fermions to scale as $g^{1/2}$, it must have either no derivatives, or both ∂_+ and ∂_- . Alternatively, it can contain one ∂_- with one left-moving and one right-moving fermions.

This difference from $AdS_5 \times S^5$ will have an important consequence. When we come to drawing Feynman diagrams in the following section, right-moving fields and derivatives will contribute powers of right-moving momenta p_- , k_- to the numerator, see (11). But the presence of left-moving objects will introduce also k_+ , and a factor $k_+ k_- = k^2$ will make any loop integral more divergent. Because of this, we will have many quadratically divergent integrals to deal with, while in the $AdS_5 \times S^5$ case [7], all of the analogous integrals were finite.

For the quartic terms, we write $\mathcal{L}_4^{\text{NFS}} = \mathcal{L}_{BB} + \mathcal{L}_{BF} + \mathcal{L}_{FF}$, and the all-boson term is the simplest:

$$\mathcal{L}_{BB} = -\frac{1}{8} [\partial_- z_i \partial_- z_i + (\partial_- y)^2 + \partial_- \bar{\omega}^\alpha \partial_- \omega_\alpha] (z_j z_j - y^2 - \frac{1}{4} \bar{\omega}^\beta \omega_\beta). \quad (18)$$

For the term mixing bosons and fermions (writing $\bar{\psi} \psi = \bar{\psi}_a \psi^a$ and $\bar{\omega} \omega = \bar{\omega}^\alpha \omega_\alpha$) we have

$$\begin{aligned} \mathcal{L}_{BF} = & -\frac{i}{8} (s_-)_a^\alpha \partial_- (s_-)_\alpha^a \left(z_i^2 - y^2 - \frac{1}{2} \omega \bar{\omega} \right) + \frac{i}{16} (s_-)_a^\alpha (s_-)_\gamma^a \omega_\alpha \overleftrightarrow{\partial}_- \bar{\omega}^\gamma - \frac{i}{32} \bar{\psi}_- \overleftrightarrow{\partial}_- \psi_- \omega \bar{\omega} \\ & + \frac{i}{8} (\bar{\psi}_- \psi_- \partial_- \omega \bar{\omega} + \bar{\psi}_- \partial_- \psi_- \omega \bar{\omega} - \bar{\psi}_- \psi_- \omega \partial_- \bar{\omega} - \partial_- \bar{\psi}_- \psi_- \omega \bar{\omega}) \\ & - \frac{i}{8} \bar{\psi}_{-a} \psi_-^b Z_c^a \overleftrightarrow{\partial}_- Z_b^c + \frac{1}{8} y [(s_-)_{a\alpha} \psi_-^a \partial_- \bar{\omega}^\alpha + (s_-)^{a\alpha} \bar{\psi}_{-a} \partial_- \omega_\alpha] \\ & - \frac{1}{16} \partial_- y [(s_-)_{a\alpha} \psi_-^a \bar{\omega}^\alpha + (s_-)^{a\alpha} \bar{\psi}_{-a} \omega_\alpha] \\ & + \frac{1}{4} y (s_-)_{a\alpha} \partial_- (s_-)_c^\alpha Z^{ca} + \frac{i}{4} (s_-)_a^\alpha \partial_- \bar{\psi}_{-c} \omega_\alpha Z^{ca} + \frac{i}{4} (s_-)_{a\alpha} \partial_- \psi_-^b \bar{\omega}^\alpha Z_b^a \\ & + \frac{i}{8} \partial_- (s_-)_a^\alpha \bar{\psi}_{-c} \omega_\alpha Z^{ca} + \frac{i}{8} \partial_- (s_-)_{a\alpha} \psi_-^b \bar{\omega}^\alpha Z_b^a \\ & - \frac{i}{8} y^2 \bar{\psi}_- \overleftrightarrow{\partial}_- \psi_- + \frac{i}{8} (s_-)_{a\alpha} (s_-)_b^\alpha Z_d^a Z'^{db} \end{aligned}$$

$$-\frac{1}{8}(\bar{\psi}_- \overset{\leftrightarrow}{\partial}_+ \psi_- + \bar{\psi}_+ \psi_- + \bar{\psi}_- \psi_+)[(\partial_- y)^2 + (\partial_- z_i)^2 + \partial_- \omega_\alpha \partial_- \bar{\omega}^\alpha]. \quad (19)$$

At this order, the powers of g under the boost (13) are the same as in the $AdS_5 \times S^5$ case [1]: terms with two right-moving fermions and ∂_- scale correctly as g . On the last line we do also have some terms with left-moving objects.

It is possible to remove the terms with left-moving factors from \mathcal{L}_{BF} by nonlinear redefinitions of the fermions, in a way which does not change final results. Similar terms were sometimes present in the $AdS_5 \times S^5$ case, in certain parameterizations of the coset⁵. We write the last term \mathcal{L}_{FF} in a form obtained by performing such a redefinition, since the result is much more compact:

$$\begin{aligned} \mathcal{L}_{FF} = & \frac{3}{8}(\bar{\psi}_- \psi_-)^2 - \frac{1}{4}\bar{\psi}_- \psi_- (\partial_+ \bar{\psi}_- \partial_- \psi_- + \partial_- \bar{\psi}_- \partial_+ \psi_-) - \frac{3}{8}\partial_+ (\bar{\psi}_- \psi_-) \partial_- (\bar{\psi}_- \psi_-) \\ & - \frac{1}{24}(s_-)^\beta_\alpha (s_-)^\delta_\beta (s_-)^\gamma_\delta (s_-)^\alpha_\gamma - \frac{1}{32}(s_-)^\alpha_\alpha (s_-)^\alpha_c \bar{\psi}_- \psi_-^c \\ & + \frac{1}{8}[\partial_+ (s_-)^\alpha_\alpha (s_-)^\alpha_c \bar{\psi}_- \partial_- \psi_-^c + \frac{1}{2}\partial_+ (s_-)^\alpha_\alpha (s_-)^\alpha_c \partial_- \bar{\psi}_- \psi_-^c \\ & + \frac{1}{2}\partial_+ (s_-)^\alpha_\alpha \partial_- (s_-)^\alpha_c \bar{\psi}_- \psi_-^c + \frac{1}{2}(s_-)^\alpha_\alpha \partial_+ (s_-)^\alpha_c \bar{\psi}_- \partial_- \psi_-^c \\ & + (s_-)^\alpha_\alpha \partial_+ (s_-)^\alpha_c \partial_- \bar{\psi}_- \psi_-^c + \frac{1}{2}\partial_- (s_-)^\alpha_\alpha \partial_+ (s_-)^\alpha_c \bar{\psi}_- \psi_-^c \\ & + \frac{1}{2}(s_-)^\alpha_\alpha (s_-)^\alpha_c \partial_- \bar{\psi}_- \partial_+ \psi_-^c + \frac{1}{2}(s_-)^\alpha_\alpha (s_-)^\alpha_c \partial_+ \bar{\psi}_- \partial_- \psi_-^c \\ & + \frac{3}{2}(s_-)^\alpha_\alpha \partial_- (s_-)^\alpha_c \bar{\psi}_- \partial_+ \psi_-^c + \frac{3}{2}\partial_- (s_-)^\alpha_\alpha (s_-)^\alpha_c \partial_+ \bar{\psi}_- \psi_-^c]. \end{aligned} \quad (20)$$

We discuss details of the redefinitions leading to this form in appendix B. Note however that this is only possible for the highest order interactions. A similar procedure for the cubic interaction \mathcal{L}_3 is not allowed.

In the following three sections we will use the above Lagrangian to calculate two-point functions on the worldsheet. We return to the full BMN theory in section 6.

3. Corrections to the light propagators

For one light mode, the exact dispersion relation is (2)

$$E = \sqrt{\frac{1}{4} + 4h(\lambda)^2 \sin^2 \frac{p_{\text{chain}}}{2}},$$

where the interpolating function is $h(\lambda) = \sqrt{\lambda/2} + c + \mathcal{O}(1/\sqrt{\lambda})$. The momentum here is the one canonically normalized for the spin chain, and it is this which scales as $p_{\text{chain}} \sim \lambda^{-1/4}$ in the near-flat-space limit, (1). Taking this into account, we can expand E^2 in λ as follows:

$$\begin{aligned} E^2 &= \frac{\lambda p_{\text{chain}}^2}{2} + \left[\frac{1}{4} + \sqrt{2\lambda} c p_{\text{chain}}^2 - \frac{\lambda p_{\text{chain}}^4}{24} \right] + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right) \\ &= p_1^2 + \frac{1}{4} + \left(\frac{4c p_1^2}{\sqrt{2\lambda}} - \frac{p_1^4}{6\lambda} \right) + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right), \end{aligned}$$

where $p_1 = \sqrt{\lambda/2} p_{\text{chain}} \sim \lambda^{1/4}$. On the second line we write it in terms of p_1 normalized to match $p_0 = E$. This p_1 is the worldsheet momentum in our uniform light-cone gauge, and its relationship to p_{chain} could in principle be deduced from the gauge but we are content just to read it off here.

In terms of the worldsheet momenta, the near-flat-space limit assumes that $p_- \gg p_+$ (or more strictly $p_- \sim \lambda^{1/4}$ and $p_+ \sim \lambda^{-1/4}$), and thus we can write $(p_1)^2 = \frac{1}{4}(p_+ - p_-)^2 =$

⁵ This is presented in [28]. For similar redefinitions, see [29, 30].

$\frac{1}{4}p_- + \mathcal{O}(1)$. Doing this in the correction terms, we arrive at

$$p_0^2 - p_1^2 = \frac{1}{4} + \left(\frac{cp_-^2}{\sqrt{2\lambda}} - \frac{p_-^4}{96\lambda} \right) + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right). \quad (21)$$

We can now read this as giving a mass correction: $p^2 = m^2 + \delta m^2$, with $m^2 = \frac{1}{4}$, and δm^2 in brackets. The *second* term there is the same two-loop correction as was calculated by [7] in the $AdS_5 \times S^5$ case⁶. We aim to compute the *first* term of the correction, containing c . The result of [2, 17] which we would like to match is that

$$c = -\frac{\log 2}{2\pi}.$$

Note that in expansion (21) the mass squared and its one- and two-loop corrections are all of the same order in $1/\sqrt{\lambda}$. This is a peculiarity of the near-flat-space limit. In the BMN case, $p_{\text{chain}} \sim \lambda^{-1/2}$ and thus $p_1 \sim \lambda^0$, restoring the natural-looking separation of orders.

Our calculation of the mass shift comes from writing a familiar geometric series involving amputated diagrams we call \mathcal{A} :⁷

$$G_\omega(p) = \frac{2i}{p^2 - \frac{1}{4}} \sum_{n=0}^{\infty} \left[\gamma \mathcal{A} \frac{2i}{p^2 - \frac{1}{4}} \right]^n$$

$$= \frac{2i}{p^2 - \left(\frac{1}{4} + \delta m^2\right)}, \quad \text{with} \quad \delta m^2 = 2i \gamma \mathcal{A}. \quad (22)$$

We will consider corrections not only to the light boson propagator $\langle \bar{\omega} \omega \rangle$ as written here, but also (in section 3.4) to all four of the light fermion propagators (11). When correcting for instance $\langle \bar{\psi}_+ \psi_+ \rangle$, the lines we amputate will no longer have fixed numerator 2 i but instead the various D_{+s} and D_{r-} , so (normalizing to the bosonic case) we absorb all of these into \mathcal{A} .

There are two topologies of diagrams we must compute, both of which contain divergences, and it is simplest to employ dimensional regularization. We show quite a lot of detail partly in order to attach later discussions about other possible cutoffs, in section 4.

3.1. Bubble diagrams for $\langle \bar{\omega} \omega \rangle$

The loop in each bubble diagram (for a light mode) always contains one light and one heavy mode. We begin with the simplest case, in which these are both bosons:

$$\mathcal{A}_B = \text{diagram} \quad , \quad q_\mu = p_\mu - k_\mu$$

$$= -1 \int \frac{d^2 k}{(2\pi)^2} \frac{B}{(k^2 - \frac{1}{4})(q^2 - 1)}, \quad B = -2 \left(\frac{p_- + k_-}{8} \right)^2.$$

The factor B includes both factors from the propagators (10) and from the vertex (17). The decorations ∂_{\pm} pull down $i k_{\pm}$ from the momentum flowing into the vertex on that line.

⁶ Of course, the conventional factors of 2 and π in (2) are different.

⁷ Recall that the interaction terms in (16) are $\sqrt{\gamma}\mathcal{L}_3 + \gamma\mathcal{L}_4$, with $\gamma = 1/g = \sqrt{2/\lambda}$.

To treat this integral, we introduce a Feynman parameter x as follows:

$$\begin{aligned}\mathcal{A}_B &= - \int \frac{d^2 k}{(2\pi)^2} \int_0^1 dx \frac{B}{\left[x \left(k^2 - \frac{1}{4} \right) + (1-x) \left((p-k)^2 - 1 \right) \right]^2} \\ &= - \int_0^1 dx \int \frac{d^2 \ell}{(2\pi)^2} \frac{B}{\left[\ell^2 - \Delta(x) \right]^2},\end{aligned}$$

where

$$\ell_\mu = k_\mu - (1-x)p_\mu, \quad \Delta(x) = \frac{1}{4}x^2 - x + 1.$$

This effective mass squared $\Delta(x)$ assumes $p^2 = \frac{1}{4}$, i.e. that we are very near to the pole of the original propagator. In terms of ℓ_μ ,

$$B = -\frac{(x-2)^2}{32}p_-^2 + \frac{2(x-2)\ell_-p_- - \ell_-^2}{32}.$$

We rotate the ℓ_0 contour to the imaginary axis, and in the resulting Euclidean space, the terms ℓ_-^n will vanish⁸. The radial integral then gives $1/2\Delta(x)$, and finally we obtain

$$\mathcal{A}_B = \frac{i}{32\pi} p_-^2 \quad \Rightarrow \quad \delta m^2 = 2i\gamma \mathcal{A}_B = \frac{-1}{16\pi} \gamma p_-^2. \quad (23)$$

Next we treat the case where the heavy and light modes in the loop are both fermions:

$$\begin{aligned} \mathcal{A}_F(p) &= \text{---}\overset{\psi^a(k)}{\circlearrowleft}_{\omega_\alpha(p)}\text{---} \\ &= 2 \int \frac{d^2 k}{(2\pi)^2} \frac{B}{(k^2 - \tfrac{1}{4})(q^2 - 1)} \end{aligned} \quad (24)$$

Here, B has the same meaning⁹, but is much more complicated since, expanding $(\mathcal{L}_3)^2$, there are in all 25 terms, and the propagators (11) give various powers of the momenta. After writing B in terms of $\ell_+^s \ell_-^{s'}$, terms with $s \neq s'$ still vanish, but there are now also divergent terms, with one or two powers of $\ell_+ \ell_- = \ell^2$ in the numerator.

As we noted below (17), the fact that left-moving momenta k_+ and q_+ appear in B , and produce divergences, is a crucial qualitative difference from the $AdS_5 \times S^5$ case.

We will treat the resulting divergent integrals using dimensional regularization. Here is the generic integral, including the case $n \neq 2$ for later use:¹⁰

$$\begin{aligned}
I_n^s(\Delta) &= \int \frac{d^d \ell}{(2\pi)^d} \frac{(\ell_+ \ell_-)^s}{[\ell^2 - \Delta]^n}, \quad d = 2 - \epsilon \\
&= \frac{i}{(2\pi)^d} \frac{2\pi^{d/2}}{\Gamma(\frac{d}{2})} \int_0^\infty dL \frac{(-1)^s L^{2s+d-1}}{[L^2 + \Delta]^n} \\
&= \frac{i}{(4\pi)^{d/2}} \frac{(-1)^s}{\Gamma(\frac{d}{2})} \frac{2}{\Gamma(\frac{d}{2})} \left(\frac{1}{\Delta}\right)^{n-s-d/2} \frac{\Gamma(n-s-\frac{d}{2})\Gamma(s+\frac{d}{2})}{2\Gamma(n)}. \quad (25)
\end{aligned}$$

⁸ We define $\ell_0 = i L_0$, $\ell_1 = L_1$ and $L_\mu = L(\sin \theta, \cos \theta)$. Then the denominator is $[L^2 - \Delta(x)]^2$, and in the numerator, any nonzero power ℓ_-^n will lead to $\int_0^{2\pi} d\theta e^{-in\theta} = 0$.

⁹ Somewhat arbitrarily, the factors from the fermion loop (i.e. re-ordering fields) and the \sum_a in the loop are part of the prefactor, not B .

¹⁰ Here of course $2\pi^{d/2}/\Gamma(\frac{d}{2})$ is the volume of a unit $(d-1)$ -sphere, and $\gamma_E = 0.577\dots$

Expanding in ϵ for $n = 2$,

$$\begin{aligned} I_2^0(\Delta) &= \frac{i}{2\pi} \frac{1}{2\Delta} \\ I_2^1(\Delta) &= \frac{i}{2\pi} \left(-\frac{1}{\epsilon} + \frac{\gamma_E - \log \pi - 2 \log 2}{2} + \frac{1}{2} + \frac{\log \Delta}{2} \right) + \mathcal{O}(\epsilon) \\ I_2^2(\Delta) &= \frac{i}{2\pi} 2\Delta \left(-\frac{1}{\epsilon} + \frac{\gamma_E - \log \pi - 2 \log 2}{2} + \frac{1}{4} + \frac{\log \Delta}{2} \right) + \mathcal{O}(\epsilon). \end{aligned}$$

(See appendix C for comments on these expansions.)

The coefficients of these integrals are as follows, after using $p_+p_- = \frac{1}{4}$:

$$\begin{aligned} B_{00} &= -\frac{3x^4 - 6x^3 - 8x^2 + 20x - 8}{32} p_-^2 \\ B_{11} &= -\frac{19x^2 - 22x}{16} p_-^2 \\ B_{22} &= -\frac{1}{4} p_-^2. \end{aligned}$$

After integrating on x , the log divergence $s = 1$ will cancel against the quadratic divergence $s = 2$. Then the final result for the fermion bubble is

$$\mathcal{A}_F = \frac{+i}{16\pi} p_-^2 \quad \Rightarrow \quad \delta m^2 = \frac{-1}{8\pi} \gamma p_-^2. \quad (26)$$

3.2. Tadpole diagrams for $\langle \bar{\omega} \omega \rangle$

The tadpoles made with two 3-vertices can be seen to vanish by looking at \mathcal{L}_3 , (17):

A diagram showing a vertex (represented by a black dot) with two external lines (dashed lines with arrows) and a loop (a circle with an arrow) attached to it. The vertex is connected to the loop by a vertical line. The external lines are horizontal and point away from the vertex. The loop is a circle with an arrow indicating a clockwise direction.

But the ones constructed with one 4-vertex are more interesting. We use (19) to calculate the first two diagrams. The only nonzero terms are those where both decorations ∂_- act on the external legs, which for ω running in the loop gives

$$\begin{aligned} \mathcal{A}_{T1} &= \begin{array}{c} \omega_\beta(k) \\ \text{---} \bullet \text{---} \\ \omega_\alpha(p) \end{array} = i \frac{1}{32} 2 p_-^2 \int \frac{d^{d=2-\epsilon} k}{(2\pi)^2} \frac{2i}{k^2 - \frac{1}{4}} \\ &= \frac{-p_-^2}{8} I_{n=1}^{s=0} (\Delta = \frac{1}{4}). \end{aligned}$$

For a heavy mode running in the loop, we obtain

$$\begin{aligned} \mathcal{A}_{T_2} &= \text{diagram of a tadpole loop with external momentum } y(k) \text{ or } z_i(k) \\ &\quad \omega_\alpha(p) = i \frac{-1}{8} (3-1) p_-^2 \int \frac{d^2 k}{(2\pi)^2} \frac{i}{k^2 - 1} \\ &= \frac{p_-^2}{4} I_1^0(1). \end{aligned} \tag{27}$$

There are also diagrams of this topology with a light fermion running in the loop. The terms in (19) which contribute are those on the last line, and they give the following two integrals

(including the only quadratically divergent tadpole):

$$\begin{aligned} \mathcal{A}_{T3} &= \text{Diagram: a tadpole with a fermion loop and a boson line} \\ &= i \frac{-i}{8} 2p_-^2 \int \frac{d^2k}{(2\pi)^2} \frac{2k_+ k_-}{k^2 - \frac{1}{4}} + i \frac{-1}{8} 4p_-^2 \int \frac{d^2k}{(2\pi)^2} \frac{-i/2}{k^2 - \frac{1}{4}} \\ &= \frac{p_-^2}{2} I_1^1\left(\frac{1}{4}\right) - \frac{p_-^2}{4} I_1^0\left(\frac{1}{4}\right). \end{aligned}$$

The diagrams with a heavy fermion in the loop can be seen to vanish from (19).

All of the tadpole integrals are $n = 1$ cases of (25) above, with $\Delta = \frac{1}{4}$ or 1. The expansions needed are

$$\begin{aligned} I_{n=1}^{s=0}(\Delta) &= \frac{i}{2\pi} \left(-\frac{1}{\epsilon} + \frac{\gamma_E - \log \pi - 2 \log 2 + \log \Delta}{2} \right) + \mathcal{O}(\epsilon) \\ I_{n=1}^{s=1}(\Delta) &= \frac{i}{2\pi} \left(-\frac{\Delta}{\epsilon} + \Delta \frac{\gamma_E - \log \pi - 2 \log 2 + \log \Delta}{2} \right) + \mathcal{O}(\epsilon). \end{aligned} \quad (28)$$

Adding the bosonic and fermionic terms, the divergences cancel leaving a total tadpole contribution of

$$\mathcal{A}_T = \frac{i}{4} p_-^2 \frac{\log 2}{2\pi} \Rightarrow \delta m_T^2 = -\frac{1}{2} \gamma p_-^2 \frac{\log 2}{2\pi}. \quad (29)$$

3.3. Total for the light boson

Adding up all of these terms, the total amplitude is $\mathcal{A}_B + \mathcal{A}_F + \mathcal{A}_T = \frac{i}{4} p_-^2 \frac{\log 2}{2\pi} + \frac{3i}{32\pi} p_-^2$. As a mass correction this is

$$\delta m^2 = -\frac{\log 2}{2\pi} \frac{\gamma}{2} p_-^2 - \frac{3}{16\pi} \gamma p_-^2. \quad (30)$$

Comparing this to (21), we see that the term with $\log 2$ (from the tadpoles) is perfect so long as $\gamma = \sqrt{2/\lambda}$. The extra term is perhaps unexpected, and we comment on this in the conclusions.

3.4. Corrections for the light fermion $\langle \bar{\psi} \psi \rangle$

We now turn to the case of corrections to the light fermion propagators (11). The two bubble diagrams we need are

$$\begin{aligned} \mathcal{A}_\psi &= \text{Diagram 1: fermion line with a boson bubble} + \text{Diagram 2: fermion line with a fermion bubble} \\ &= \int \frac{d^2k}{(2\pi)^2} \frac{s_\alpha^a(q)}{(k^2 - \frac{1}{4})(q^2 - 1)} B. \end{aligned}$$

The integrals are of course the same as (24) above, with different coefficients $B = \sum_{s,s'} \ell_+^s \ell_-^{s'} B_{ss'}$. As noted above, B now includes the numerators $D_{\psi\psi}$ of the two amputated propagators (as well as the vertices and internal propagators). We must calculate the coefficients separately for each of the four cases $\langle \bar{\psi}_+ \psi_+ \rangle$, $\langle \bar{\psi}_- \psi_- \rangle$, $\langle \bar{\psi}_- \psi_+ \rangle$ and $\langle \bar{\psi}_+ \psi_- \rangle$, and the results are all different until the point at which we use $p^2 = \frac{1}{4}$ to get the shift near to the pole of the

unperturbed propagator. Then they all agree, and we obtain

$$\begin{aligned} B_{00} &= -\frac{1}{128}(64 + 24x - 32x^2 - 8x^3 + x^4 + 2x^5)p_-^2 \\ B_{11} &= -\frac{1}{16}(-12 - 7x + x^2 + 6x^3)p_-^2 \\ B_{22} &= -\frac{1}{8}(6x - 1)p_-^2. \end{aligned}$$

Then using the expansions of the integrals in (25), we find

$$\delta m^2 = \frac{1}{64\pi} \gamma \left(-\frac{42}{\epsilon} + 21\gamma_E - 26 \log 2 + 21 \log \pi \right) p_-^2. \quad (31)$$

There is an important observation we can make at this stage. In this expansion in $1/\epsilon$, there is no order 1 term with a rational coefficient (inside the bracket)¹¹. Since no such terms will appear in any of the tadpole integrals (28), we conclude that there will be no ‘extra’ term like that seen in (30). A further difference from the ω case above is that the total of the bubble diagrams is not finite.

Next we turn to the tadpole diagrams, using (19) and (20). We must compute these for each of the four cases $\langle \bar{\psi}_+ \psi_+ \rangle$, $\langle \bar{\psi}_- \psi_- \rangle$, $\langle \bar{\psi}_- \psi_+ \rangle$ and $\langle \bar{\psi}_+ \psi_- \rangle$ separately, but again the final coefficients work out to be the same¹². Those with a light mode in the loop are

$$\begin{aligned} \mathcal{A}_L &= \text{diagram with } \omega_\beta(k) \text{ loop and } \psi^a(p) \text{ external line} + \text{diagram with } \psi^b(k) \text{ loop and } \psi^a(p) \text{ external line} \\ &= p_-^2 \left[\frac{9}{16} I_1^0\left(\frac{1}{4}\right) - \frac{3}{8} I_1^1\left(\frac{1}{4}\right) \right]. \end{aligned}$$

With a heavy mode in the loop, we find

$$\begin{aligned} \mathcal{A}_H &= \text{diagram with } y(k) \text{ or } z_i(k) \text{ loop and } \psi^a(p) \text{ external line} + \text{diagram with } s_\beta^b(k) \text{ loop and } \psi^a(p) \text{ external line} \\ &= p_-^2 \left[-\frac{3}{2} I_1^0(1) + \frac{3}{8} I_1^1(1) \right]. \end{aligned} \quad (32)$$

After using the expansions of the integrals in (28), the total is as follows:

$$\delta m^2 = \gamma \frac{21}{64\pi} \left(-\frac{2}{\epsilon} + \gamma_E + \log 2 - 2 \log 2 \right) p_-^2.$$

Happily the divergent $1/\epsilon$ term cancels that of (31).

Adding the bubble and tadpole contributions gives us that the full mass shift, which for all the $\langle \bar{\psi} \psi \rangle$ cases is

$$\delta m^2 = -\gamma \frac{\log 2}{4\pi} p_-^2. \quad (33)$$

This is exactly what we would expect from $c = -\frac{\log 2}{2\pi}$ in (21). Unlike the bosonic case (30), there are no extra terms.

¹¹ Such terms do occur in the individual bubble diagrams drawn, which contribute $\pm \frac{3}{64\pi} p_-^2$, but these cancel out of the total.

¹² Note also that whether we use (B.1) or the simplified (20) makes no difference to these coefficients.

4. A detour via some delicate cutoffs

In this section, we look at some effects of employing explicit energy or momentum cutoffs Λ instead of dimensional regularization.

The main reasons for doing so are to try to better understand the extra terms in the mass correction (30), and to explore the prescription dependence of c by looking at ‘old’ and ‘new’ cutoffs (in section 4.3). We will also get extra cross-checks between the bubble (\mathcal{L}_3) and the tadpole (\mathcal{L}_4) calculations.

Most of this section will be about only the light boson case. We comment briefly on other cases in section 4.4.

4.1. Using hard cutoffs

The first observation is that if we work with hard momentum cutoffs $|k| < \Lambda$, instead of dimensional regularization, then the cancellation of infinities is different. Each of \mathcal{A}_F and \mathcal{A}_T contains a logarithmic and a quadratic divergence, and above we cancelled these *within* them, since all are $1/\epsilon$ terms when working in $d = 2 - \epsilon$ dimensions. However in terms of a hard cutoff, instead these Λ^2 terms cancel *between* them. This cancellation provides a useful cross-check between the bubble (\mathcal{L}_3) and the tadpole (\mathcal{L}_4) calculations.

The only quadratically divergent terms are the $(\ell_+ \ell_-)^2$ part of the fermionic bubble integral¹³

$$\begin{aligned}\mathcal{A}_F^{s=2} &= 2B_{22} \frac{i}{2\pi} \int_0^1 dx \int_0^\Lambda dL \frac{L^5}{[L^2 + \Delta(x)]^2} \\ &= \frac{i p_-^2}{4\pi} \left[\frac{\Lambda^2}{2} - \frac{7}{6} \log \Lambda + \frac{\log 2}{6} - \frac{7}{72} + \mathcal{O}\left(\frac{1}{\Lambda}\right) \right]\end{aligned}\quad (34)$$

and the fermionic tadpole integral

$$\begin{aligned}\mathcal{A}_{T3} &= \frac{p_-^2}{2} \frac{i}{2\pi} \int_0^\Lambda dK \frac{K^3}{K^2 + \frac{1}{4}} - \frac{p_-^2}{4} \frac{i}{2\pi} \int_0^\Lambda dK \frac{-K}{K^2 + \frac{1}{4}} \\ &= \frac{i p_-^2}{4\pi} \left[\frac{\Lambda^2}{2} - \frac{1}{4} \log \Lambda - \frac{\log 2}{4} \right] + \frac{i p_-^2}{4\pi} [2 \log \Lambda + 2 \log 2] + \mathcal{O}\left(\frac{1}{\Lambda}\right).\end{aligned}\quad (35)$$

The $\log \Lambda$ divergences similarly cancel. They are

$$\begin{aligned}\mathcal{A}_F^{s=1} &= 2 \int_0^1 dx B_{11} \int_0^\Lambda dL \frac{-L^3}{[L^2 + \Delta(x)]^2} \\ &= \frac{i p_-^2}{4\pi} \left[-\frac{7}{6} \log \Lambda - \frac{17 \log 2}{6} + \frac{155}{72} + \dots \right] \\ \mathcal{A}_{T1} &= \frac{-p_-^2}{8} \int_0^\Lambda dK \frac{-K}{K^2 + \frac{1}{4}} = \frac{i p_-^2}{4\pi} \left[\frac{1}{4} \log \Lambda + \frac{1}{4} \log 2 + \dots \right] \\ \mathcal{A}_{T2} &= \frac{p_-^2}{4} \frac{i}{2\pi} \int_0^\Lambda dK \frac{-K}{K^2 + 1} = \frac{i p_-^2}{4\pi} \left[-\frac{1}{2} \log \Lambda + \dots \right]\end{aligned}\quad (36)$$

and the finite terms are of course unchanged:

$$\mathcal{A}_F^{s=0} = \int_0^1 dx 2B_{00} \frac{1}{2\Delta(x)} = \frac{i p_-^2}{4\pi} [3 \log 2 - 2]$$

¹³ We note in passing the following issue [31]. In four dimensions, the shift from integrating over k_μ with $|k| < \Lambda$ to integrating over ℓ_μ with $|\ell| < \Lambda$ would change the value of the Λ^2 -divergent integral, and thus not be allowed. (The finite and $\log \Lambda$ integrals are unchanged.) This is however not a problem in two dimensions, nor is it a problem in dimensional regularization, in any number of dimensions.

$$\mathcal{A}_B = \frac{i p_-^2}{4\pi} \frac{1}{8}.$$

Adding these up, we obtain the same finite result as before (30):

$$\delta m^2 = -\frac{\log 2}{2\pi} \frac{\gamma}{2} p_-^2 - \frac{3}{16\pi} \gamma p_-^2.$$

Note in particular that we have exactly the same ‘extra’ term as we had using dimensional regularization. We now turn to looking for other explanations of this term, for this case.

4.2. High-precision cutoffs

Note that the finite part after the cancellation of quadratic divergences (between bubble and tadpole) is sensitive to very small changes in the cutoff. In this section, we investigate the effects of such small changes, replacing Λ with one of these:

$$\begin{aligned} \Lambda_{\text{light}} &= \Lambda - \frac{\iota}{2\Lambda}, & \text{light modes} \\ \Lambda_{\text{heavy}} &= \Lambda - \frac{\kappa}{2\Lambda}, & \text{heavy modes.} \end{aligned} \quad (37)$$

For the next page or so we leave ι and κ arbitrary.

- For the tadpole we can simply write down the result, since the only Λ^2 divergent integral (35) involves only a light mode. The effect of changing from $\Lambda \rightarrow \Lambda_{\text{light}}$ is 1 i:

$$\Lambda^2 \rightarrow \Lambda^2 - \iota. \quad (38)$$

- For the bubble’s $s = 2$ term (34), we expect some change involving both heavy and light modes, but it is not obvious what mixture of ι and κ should appear. In fact, it is not clear that whether there will be any way of implementing *both* cutoffs (37) in the same loop integral. But we can make an attempt by introducing Pauli–Villars regulators [32], replacing in the heavy and light propagators the following:

$$\begin{aligned} \frac{1}{k^2 - \frac{1}{4}} &\xrightarrow{\text{PV}} \frac{1}{k^2 - \frac{1}{4}} - \frac{1}{k^2 - \Lambda_{\text{light}}^2} \\ \frac{1}{q^2 - 1} &\xrightarrow{\text{PV}} \frac{1}{q^2 - 1} - \frac{1}{q^2 - \Lambda_{\text{heavy}}^2}. \end{aligned}$$

The propagators are unchanged for $k^2 \ll \Lambda^2$, but when $k^2 \gg \Lambda^2$ they die like $1/k^4$. Defining a generalization of the integral in (24) to allow two arbitrary masses

$$J(m, M) = \int \frac{d^2 k}{(2\pi)^2} \frac{B}{(k^2 - m^2)(q^2 - M^2)}$$

the effect of imposing the Pauli–Villars regulators is to replace $J(\frac{1}{2}, 1)$ with four terms:

$$J(\tfrac{1}{2}, 1) \xrightarrow{\text{PV}} J(\tfrac{1}{2}, 1) - J(\Lambda_{\text{light}}, 1) - J(\tfrac{1}{2}, \Lambda_{\text{heavy}}) + J(\Lambda_{\text{light}}, \Lambda_{\text{heavy}}).$$

Each of these terms can be treated exactly as we did before¹⁴, simply re-using the integrals (25) with the appropriate effective masses $\Delta(x, m, M)$. Thus the $s = 2$ part of each is given by

$$J^{s=2}(m, M) = \int_0^1 dx \int \frac{d^{2-\epsilon} \ell}{(2\pi)^2} \frac{(\ell_+ \ell_-)^2}{[\ell^2 - \Delta(x, m, M)]^2},$$

¹⁴ This means that we are regulating each of the four terms with dimensional regularization. We can instead use another hard cutoff Λ' for all the terms, and as long as $\Lambda' \gg \Lambda_{\text{heavy}}, \Lambda_{\text{light}}$, the result is identical. (If $\Lambda' = \Lambda$, then (40) becomes $\Lambda^2 \rightarrow \Lambda^2 - (\iota + \kappa)$ instead.)

where

$$\begin{aligned}\ell_\mu &= k_\mu - (1-x)p_\mu \\ \Delta &= xm^2 + (1-x)M^2 + x(1-x)p^2.\end{aligned}$$

Note that ℓ_μ is defined in the same way in all four terms, so that the identification of the term $(\ell_+\ell_-)^2$ in B is the same here as before. (It comes with the same coefficient B_{22} .) The final result for the $s = 2$ integral (at $p^2 = \frac{1}{4}$) is then

$$J_{\text{PV}}^{s=2}\left(\frac{1}{2}, 1\right) = \frac{\Lambda^2}{2} - \frac{7}{6} \log \Lambda + \left(\frac{\log 2}{6} - \frac{5}{48} - \frac{\iota + \kappa}{4}\right) + \mathcal{O}\left(\frac{1}{\Lambda}\right). \quad (39)$$

Thus, the effect on the bubble diagram of turning on ι, κ can be summarized as

$$\Lambda^2 \rightarrow \Lambda^2 - \frac{\iota + \kappa}{2}. \quad (40)$$

This makes some sense, in that if you change both cutoffs by the same amount $\iota = \kappa$ then this becomes an overall change of Λ^2 . What you cannot predict without calculation is that the effect of $\iota \neq \kappa$ should not be some other mixture.

The total effect on \mathcal{A} of both (38) and (40) is then as follows. Using (34) and (35) above, we have $\mathcal{A} \rightarrow \mathcal{A} + \frac{i p_-^2}{8\pi} \left(\iota - \frac{\iota + \kappa}{2}\right)$, or, in terms of the mass shift,

$$\delta m^2 = 2 i \mathcal{A} \rightarrow \delta m^2 + (\kappa - \iota) \frac{\gamma p_-^2}{8\pi}.$$

It remains to argue what tiny shifts ι, κ we should use.

Ideally, one would like to impose exactly the same cutoff on the physical energy of all the modes. Unlike the worldsheet momentum k_1 , or worse the Euclidean energy K_0 , the energy is a physical, gauge-invariant quantity.

One argument is this: in the original Lorentzian momentum, $E^2 = k_0^2 = k_1^2 + m^2$ on the mass shell. Imposing $|E| < \Lambda$ thus reads $k_1^2 < \Lambda^2 - m^2$. If we impose this momentum cutoff on both of the Euclidean directions, then we are led to $K^2 < \Lambda^2 - m^2$. For the light modes, this means $\iota = \frac{1}{4}$, while for the heavy modes $\kappa = 1$. This gives

$$\delta m^2 \rightarrow \delta m^2 + \frac{3}{32} \frac{p_-^2 \gamma}{\pi}. \quad (41)$$

Sadly this is too small to cancel the extra term in (30), by a factor of 2.

The attentive reader will by this point have smelled something a little fishy. Not only have we made up a fairly strange cutoff, (37), we have also mixed the results of using this in the Pauli–Villars bubble integral (39) with that of using it in a tadpole integral calculated with a hard momentum cutoff (35). The reason for doing so is that Pauli–Villars is not strong enough to regulate the tadpole integral, and we do not know of any regulator both strong enough and potentially sensitive to independent variations of Λ_{heavy} and Λ_{light} . Nevertheless, the complaint is well founded; note that one of the finite terms from Pauli–Villars differs from that given by the hard cutoff—compare (39) to (34).

What we believe we *have* shown is that justifiable alterations to the cutoffs used alter δm^2 by terms of the form $\gamma p_-^1/\pi \times (\text{rational})$, which is the same shape as the extra term in (30). And also that these do *not* produce terms of the form $p_-^2 \log 2/\pi$. That the cancellation is not perfect is perhaps a limitation of our implementation of this physical cutoff.

4.4. Modes other than the light boson

So far we have discussed in section 4 only the ω particle. Let us first comment on how the issues of section 4.3 translate to other cases.

- For the heavy bosons which we treat in the following section, the observation that the new sum sets the tadpoles to zero carries over trivially; the calculation of the tadpole contribution is identical. However, the bubble diagram contains two light modes; thus our objection to the new sum does not hold for these.
- For the light fermion ψ , the tadpole (32) also contains the Λ^2 -divergent integral $I_1^1(1)$, for which

$$I_1^1(1) \xrightarrow{\text{new}} 4 I_1^1\left(\frac{1}{4}\right) = I_1^1(1) + \frac{i}{2\pi} \frac{\log \frac{1}{4}}{2}.$$

This change is identical to that for $I_1^0(1)$ above. Then it is easy to see that while we do not create a divergence, the finite change (from trying to go to the new sum) is *not* so simple in this case, i.e. it does not cancel the $\log 2$ term in (33).

The fact that we cannot cancel the $\log 2$ like this for ψ is possibly related to another issue. If we attempt to do the whole calculation with a momentum cutoff Λ (generalizing section 4.1), then we do not get a finite result. The reason for this is not entirely clear to us, but let us observe here that the cancellations of $1/\epsilon$ terms which make the result using dimensional regularization finite are highly nontrivial—see (33), and (47) below.

This is true also for the heavy bosons of the function section, where for instance it is clear that the bubble \mathcal{A}_b is finite (44), but the tadpoles \mathcal{A}_T are as for the ω case, and thus Λ^2 divergent (35).

5. Correction to the heavy boson propagators

For a heavy mode, we should start with the following dispersion relation:

$$E_{\text{heavy}} = \sqrt{1 + 16h(\lambda)^2 \sin^2 \frac{p_{\text{chain}}}{4}} = 2E_{\text{light}}\left(\frac{p_{\text{chain}}}{2}\right).$$

This is simply the energy of two superimposed light modes, written in terms of their *total* momentum. This relation was confirmed to hold at one loop in the giant magnon regime $p_{\text{chain}} \sim 1$ in [17].¹⁶ Expanding E^2 in $1/\sqrt{\lambda}$ exactly as for the light mode case, and in particular using the same normalization $p_1 = \sqrt{\lambda/2} p_{\text{chain}}$, we obtain

$$p_0^2 - p_1^2 = 1 + \left(\frac{cp_-^2}{\sqrt{2\lambda}} - \frac{p_-^4}{384\lambda} \right) + \mathcal{O}\left(\frac{1}{\sqrt{\lambda}}\right). \quad (43)$$

The mass correction δm^2 is again the term in brackets. Its first term, which we aim to compute here, should be identical to that for the light modes.

The calculation of course uses a similar geometric series to (22), although this time $\delta m^2 = i \gamma \mathcal{A}$, since y and z_i (unlike ω^α) are canonically normalized.

After working out the corrections for fields y and z_i we discuss the issue of their stability in section 5.3.

¹⁶ This giant magnon, in an RP^2 subspace, is a nonlinear superposition of two elementary magnons, as was shown by [34]. When discussing giant magnons it is often convenient to write $E = \sqrt{1 + 8\lambda \sin^2(p'/2)}$, where p' is the momentum of one of the constituent magnons, not the total. (See for instance section 4.3 of [35].)

5.1. Diagrams correcting $\langle yy \rangle$

Recall that for the light boson, the bosonic tadpoles came from (18) and the fermionic ones from the last line of (19), which is

$$\mathcal{L}_{\text{tad}} = \frac{-1}{8} [\partial_- z_i \partial_- z_i + (\partial_- y)^2 + \partial_- \bar{\omega}^\alpha \partial_- \omega_\alpha] (i \bar{\psi}_{-b} \overset{\leftrightarrow}{\partial}_+ \psi_-^b + \bar{\psi}_{-b} \psi_+^b + \bar{\psi}_{+b} \psi_-^b).$$

Both of these terms share the same first factor, and by looking at this factor it is clear that we can re-use exactly the same diagrams when the external particle is y . The only change is a factor of 2 because y is real, while ω_α was complex, but since we now have $\delta m^2 = i \gamma \mathcal{A}$ (without a 2), the correction δm^2 is unchanged:

$$\begin{aligned} \mathcal{A}_{T'} &= \text{diagram with } \omega_\alpha \text{ or } \psi^a \text{ in loop} + \text{diagram with } y \text{ or } z_i \text{ in loop} = 2\mathcal{A}_T \\ &= \frac{i}{2} p_-^2 \frac{\log 2}{2\pi} \\ &\Rightarrow \delta m_T^2 = -\frac{1}{2} \gamma p_-^2 \frac{\log 2}{2\pi} \quad \text{the same as (29).} \end{aligned}$$

There is only one bubble diagram to draw, which has two ω_α particles in the loop. The contribution is as follows:

$$\begin{aligned} \mathcal{A}_y &= \text{diagram with } \omega_\alpha(k) \text{ and } \omega_\alpha(q) \text{ in loop} \quad , \quad \text{now with } q_\mu = k_\mu - p_\mu \\ &= i^2 2 \int \frac{d^2 k}{(2\pi)^2} \frac{2i}{(k^2 - \frac{1}{4})} \frac{2i}{(q^2 - \frac{1}{4})} \left(\frac{k_- + q_-}{8} \right)^2 \\ &= \frac{1}{8} \int_0^1 dx \int \frac{d^2 \ell}{(2\pi)^2} \frac{(2\ell_- + (1-2x)p_-)^2}{[\ell^2 - \Delta'(x)]^2} \\ &= \frac{1}{8} p_-^2 \frac{i}{2\pi} \int_0^1 dx \frac{(1-2x)^2}{2\Delta'(x)} \\ &= \frac{i}{8\pi} p_-^2. \end{aligned} \tag{44}$$

where $\Delta'(x) = x^2 - x + \frac{1}{4} = \frac{1}{4} \Delta(4x)$ and (as before) $\ell = k - (1-x)p$.

Thus the total mass shift for the y particle is

$$\delta m^2 = -\frac{1}{2} \gamma p_-^2 \frac{\log 2}{2\pi} - \frac{1}{8\pi} \gamma p_-^2.$$

As for the light boson ω in (30), there is an extra term not expected from the dispersion relation (43). This now comes from the only bubble diagram possible, rather than the addition of a bosonic and a fermionic bubble, and thus it is difficult to imagine a cancellation of this term.

5.2. Diagrams correcting $\langle z_i z_j \rangle$

The tadpole corrections are completely identical to those for the $\langle yy \rangle$ case above, by the same argument used there.

The only bubble diagram has ψ fields in the loop:

The diagram shows a central vertex labeled $z_i(p)$. Two dashed lines extend horizontally from the vertex, one to the left and one to the right. A circular loop is attached to the vertex. The top arc of the loop is labeled $\psi^a(k)$ with an arrow pointing clockwise. The bottom arc of the loop is labeled $\psi^b(q)$ with an arrow pointing counter-clockwise.

Since the terms in \mathcal{L}_3 are written with Z_b^a instead of z_i , it is easiest to work in terms of these, and so we write $\mathcal{A}_z = 2\mathcal{A}_{01}^{10}$ and calculate the latter. Here is the integral:

$$\begin{aligned} \mathcal{A}_{01}^{10} &= -i^2 \int \frac{d^2 k}{(2\pi)^2} \frac{B}{(k^2 - \frac{1}{4})(q^2 - \frac{1}{4})} \\ &= \int_0^1 dx \int \frac{d^2 \ell}{(2\pi)^2} \frac{B_{00} + \ell_+ \ell_- B_{11} + (\ell_+ \ell_-)^2 B_{22}}{[\ell^2 - \Delta'(x)]^2}, \quad \Delta' = x^2 - x + \frac{1}{4}, \end{aligned}$$

where as before $B = \sum_{s,s'} \ell_+^s \ell_-^{s'} B_{ss'}$, and we use the same $\Delta'(x)$ as for the $\langle yy \rangle$ case above. After using $p_+ p_- \approx 1$, the coefficients needed are

$$\begin{aligned} B_{00} &= -\frac{1}{8} (2x-1)^2 (3x^2-3x-2) p_-^2 \\ B_{11} &= \frac{-22x^2+22x-3}{4} p_-^2 \\ B_{22} &= -p_-^2. \end{aligned}$$

Expanding, using (25), once again the logarithmic and quadratic divergences cancels within the bubble term. The final result is

$$\mathcal{A}_z = 2 \left(-\frac{i p_-^2}{8\pi} \right).$$

The total mass shift for the z particles is thus

$$\delta m^2 = -\frac{1}{2}\gamma p_-^2 \frac{\log 2}{2\pi} + \frac{1}{4\pi}\gamma p_-^2.$$

5.3. Breaking of mass coincidence

The heavy modes have $m = 1$, precisely twice the light modes' $m = \frac{1}{2}$. And the quantum numbers work out that there is a pair of light modes which carries all the same indices as each heavy mode¹⁷. One can ask how this gets modified at one loop, and we begin with some rather simple observations as follows.

(i) By expanding the dispersion relation we had that, for both heavy and light modes,

$$\delta m^2 = c \frac{p_-^2}{\sqrt{2\lambda}} + \text{two loops.}$$

This implies $2 \times \Delta m_{\text{light}} = 2\delta m^2$ and $\Delta m_{\text{heavy}} = \frac{1}{2}\delta m^2$. Using $c = -\log 2/2\pi \approx -0.11 < 0$, we conclude that after these corrections the heavy mode is *more* massive than a pair of light modes. Thus, it is kinematically allowed to decay into two light modes.

¹⁷ As can be seen from the interaction terms in (17). The decomposition of course matches that used in the algebraic curve for the construction of off-shell frequencies [36, 33, 17].

- (ii) By direct calculation, we find in addition to this some other terms. These are not large enough to alter the conclusion. Here are the numbers after including these terms:

$$2 \Delta m_\omega = -0.23 \gamma p_-^2$$

$$2 \Delta m_\psi = -0.11 \gamma p_-^2$$

and for heavy modes,

$$\Delta m_y = -0.13 \gamma p_-^2$$

$$\Delta m_z = -0.071 \gamma p_-^2.$$

The decays $y \rightarrow \omega + \bar{\omega}$ and $z \rightarrow \psi + \bar{\psi}$ are thus still allowed.

- (iii) If $c = 0$, then we expect that $\delta m^2 = 0$, and so at this order the coincidence of masses remains unbroken.

This issue was investigated in a more subtle way by Zarembo in [21], and this was continued in [13]. The idea is this: instead of working exactly at the pole of the original propagator, $p^2 = 1$ for a heavy mode, we can make an expansion in $p^2 - 1$ and look at the effect of the second term.

If we assume that $p^2 < 1$, then the following integral is real:

$$\begin{aligned} \int_0^1 dx \frac{(1-2x)^2}{\frac{1}{4} - x(1-x)p^2} &= \frac{4}{p^2} - \frac{4\sqrt{1-p^2}}{p^3} \arcsin(p) \\ &= 4 - 2\pi\sqrt{1-p^2} + 8(1-p^2) + \mathcal{O}(1-p^2)^{3/2}. \end{aligned}$$

This integral is the generalization away from $p^2 = 1$ of that appearing in the $\langle yy \rangle$ bubble (44). It comes with a factor $i/2\pi$ in \mathcal{A} and a further factor $i\gamma$ in δm^2 —in all a minus sign. Thus the term 4 here contributes to $\delta m^2 < 0$, ensuring $p^2 < 1$ self-consistently. In our full calculation there is also the tadpole contribution (proportional to $\log 2$), but this too is negative, as noted above. The conclusion is that even including the expansion in $p^2 - 1$, the mass corrections are all real, and so the pole of the corrected propagator $G_y(p)$ remains on the real axis.

While this integral diverges for real $p^2 > 1$, the expansion given does hold for complex p^2 . One might wonder whether, if something cancelled the leading term 4, the second term might amount to a complex mass correction. One way to investigate this is as follows. Define some coefficients B, C, D, F like

$$\begin{aligned} p^2 &= 1 + \delta m^2, \\ \delta m^2 &= -\gamma[B + C\sqrt{1-p^2} + D(1-p^2) + \mathcal{O}(1-p^2)^{3/2}] + \gamma^2 F + \mathcal{O}(\gamma^3). \end{aligned}$$

Then $B > 0$ is the simple case already discussed. If $B = 0$, then $C > 0$ looks like that it will guarantee a negative mass shift, thus keeping $\sqrt{1-p^2}$ real, while $C < 0$ is not obvious. But we can solve this equation explicitly for p^2 , and expanding in γ , we obtain

$$p^2 = 1 - B\gamma \pm \sqrt{BC} \gamma^{3/2} + \left(-\frac{C^2}{2} - BD + F\right) \gamma^2 + \mathcal{O}(\gamma^{5/2}).$$

Now it seems clear what happens if $B = 0$; regardless of the sign of C the correction is still a real number, but it enters only at two loops.

6. Corrections in the BMN Limit

In this section we repeat our calculations above using the BMN Lagrangian of [13]. The physical reason for doing so is that we want to check that the near-flat-space limit and the diagrammatic calculation of δm^2 commute.

When taking the near-flat-space limit of our BMN results, in fact we recover not only the term arising from $c = -\log 2/2\pi$ in the dispersion relation, but also the extra terms seen above (for each particular mode). This indicates that these extra terms cannot be artefacts of the simpler limit. In the BMN calculation, we also find some constant terms which are not visible in the near-flat-space limit, but which also break supersymmetry.

The approach and the diagrams needed are identical to those for the near-flat-space limit, but the number of terms involved is much greater. Consequently, we show much less detail here. But this also means that the cancellation of divergences is more delicate than it was, and this provides further checks on our work.

6.1. The dispersion relation

We can expand the dispersion relation (2) in the same way as for the near-flat-space case (21). The normalization of the worldsheet momentum is exactly the same $p_{\text{chain}}\sqrt{\lambda/2} = p_1$, as this is a statement only about the gauge we are in. But now we have $p_{\text{chain}} \sim \lambda^{-1/2}$ from (1), which changes the result to

$$E^2 = \frac{1}{4} + p_1^2 + \frac{2\sqrt{2}}{\sqrt{\lambda}} c p_1^2 + \mathcal{O}\left(\frac{1}{\lambda}\right). \quad (45)$$

This reads $p_0^2 - p_1^2 = \frac{1}{4} + \delta m^2$, and thus we expect to see

$$\delta m^2 = \frac{2\sqrt{2}}{\sqrt{\lambda}} c p_1^2.$$

This time, the correction term δm^2 is of order $1/\sqrt{\lambda}$, while p_0^2 and p_1^2 are of order 1. (The two-loop term we wrote in (21) is now of order $1/\lambda$.)

For a heavy mode, $E_{\text{heavy}}(p) = 2E(p/2)$, we again obtain $p_0^2 - p_1^2 = 1 + \delta m^2$ with the same δm^2 at this order.

6.2. Results for light modes

For the light bosonic propagator we find

$$\begin{aligned} \langle \bar{\omega} \omega \rangle : \quad \delta m^2/\gamma &= \frac{9}{16\pi\epsilon} - \frac{3}{4\pi} p_1^2 + \dots \quad \text{from bubble diagrams} \\ &\quad - \frac{9}{16\pi\epsilon} - \frac{\log 2}{\pi} p_1^2 + \dots \quad \text{from tadpoles} \\ &= -\frac{5}{96\pi} - \frac{3}{4\pi} p_1^2 - \frac{\log 2}{\pi} p_1^2 \quad \text{in total.} \end{aligned} \quad (46)$$

The dots for bubble and tadpole contributions are constant (and non-divergent) terms. If we now write $p_1 = \frac{1}{2}(p_+ - p_-)$ and perform the worldsheet boost, then we end up with the result obtained starting from the near-flat-space Lagrangian. Thus the limit commutes with the calculation.

However, for the BMN case, we see that there is one additional nontrivial $1/\epsilon$ cancellation between the bubbles and the tadpoles which is invisible in the near-flat-space calculation (where all $1/\epsilon$ cancelled within each diagram). This provides a further consistency check.

For the light fermionic propagator we have

$$\begin{aligned} \langle \bar{\psi}_- \psi_+ \rangle : \quad \delta m^2/\gamma &= \frac{7}{64\pi\epsilon} + \frac{1}{8\pi} \left(\frac{21}{\epsilon} - \frac{21\gamma_E}{2} + 13 \log 2 - \frac{21 \log \pi}{2} \right) p_1^2 + \dots \text{bubbles} \\ &\quad - \frac{7}{64\pi\epsilon} - \frac{1}{8\pi} \left(\frac{21}{\epsilon} - \frac{21\gamma_E}{2} + 21 \log 2 - \frac{21 \log \pi}{2} \right) p_1^2 + \dots \text{tadpoles.} \end{aligned}$$

The coefficients in this depend on which of the four cases $\langle \bar{\psi}_{\pm} \psi_{\pm'} \rangle$ we consider. But for all cases, the total is the same:

$$\langle \bar{\psi} \psi \rangle : \quad \delta m^2 / \gamma = -\frac{1}{96\pi} - \frac{\log 2}{\pi} p_1^2. \quad (47)$$

Again we see a cancellation of $1/\epsilon$ divergences in both the constant and the p_1^2 terms. And again the result reduces to the near-flat-space one (33).

Note that in both of these results there are extra constant terms. These are of course invisible in the near-flat-space limit. But like the extra terms seen there, they are not the same for the boson and the fermion, thus breaking supersymmetry.

6.3. Results for heavy bosons

Starting out with the y mode, we have

$$\begin{aligned} \langle yy \rangle : \quad \delta m^2 / \gamma &= \frac{1}{2\pi\epsilon} - \frac{1}{2\pi} p_1^2 + \dots && \text{bubbles} \\ &- \frac{1}{2\pi\epsilon} - \frac{\log 2}{\pi} p_1^2 + \dots && \text{tadpoles} \\ &= -\frac{1}{4\pi} - \frac{1}{2\pi} p_1^2 - \frac{\log 2}{\pi} p_1^2. \end{aligned} \quad (48)$$

The complex piece of δm^2 comes from the x integration in the Feynman parameterization.

Finally, for the z_i modes (transverse AdS_4 directions) we have

$$\begin{aligned} \langle z_i z_j \rangle : \quad \delta m^2 / \gamma &= \frac{1}{4\pi\epsilon} - \frac{p_1^2}{\pi} + \dots && \text{bubbles} \\ &- \frac{1}{4\pi\epsilon} - \frac{\log 2}{\pi} p_1^2 + \dots && \text{tadpoles} \\ &= -\frac{2}{3\pi} - \frac{p_1^2}{\pi} - \frac{\log 2}{\pi} p_1^2. \end{aligned} \quad (49)$$

As with the light propagators, all corrections reduce to those calculated using the near-flat-space Lagrangian. Thus, at least for two-point functions, the truncation of the BMN theory to the near-flat-space limit is one-loop quantum consistent.

7. Conclusions

The results of this paper are as follows.

- (i) We have computed one-loop corrections to propagators in both the near-flat-space and BMN limits, and we find that our calculations ‘commute with the limit’, meaning that the near-flat results are recovered as limits of the BMN ones. This is the first such comparison in $AdS_4 \times CP^3$, and is evidence that the near-flat-space limit is a consistent truncation at one loop.
- (ii) Comparing our results to the dispersion relation, we expect δm^2 to be proportional to the $c = -\log 2/2\pi$ in the expansion of $h(\lambda)$, and we correctly find this term. We also (in most cases) find some extra terms. Our results are always of the form

$$\begin{aligned} \delta m^2 &= -\frac{\log 2}{2\pi} 2\gamma p_1^2 + U \frac{\gamma p_1^2}{\pi} + V \frac{\gamma}{\pi} && \text{BMN limit} \\ &\rightarrow -\frac{\log 2}{2\pi} \frac{\gamma p_1^2}{2} + U \frac{\gamma p_1^2}{4\pi} && \text{near-flat-space,} \end{aligned} \quad (50)$$

where U and V are rational numbers depending on the mode being studied.

- (iii) The term with $\log 2$ comes from¹⁸ the tadpole diagrams, and for these diagrams, we can implement the ‘new’ summation prescription of Gromov and Mikhailov [3]. Doing so changes the result to remove this $\log 2$ term perfectly, corresponding to $c = 0$. This agrees with expectations from [3, 2, 17, 4]. However, we *cannot* make this change of prescription for the bubble diagrams, since the same loop momentum applies to both heavy and light modes.

7.1. Extra terms U and V in (50)

Perhaps the first thing to note is that these terms break supersymmetry. Most of our calculation is done using dimensional regularization, and so it is natural to suspect that this regulator is the culprit. (Supersymmetry after all is very different in different dimensions of spacetime.) It is possible that the terms U and V should be absorbed into a (finite) renormalization of the bare masses $m = \frac{1}{2}, 1$, restoring supersymmetry [37]. In this case the correct attitude towards our calculation is that we have used the exact answer for the mass shift (from the dispersion relation) to work out what bare masses must be used in the propagators for future calculations, done with the same regulator.

For the light boson, we were able to repeat our calculation using instead an explicit momentum cutoff (in section 4.1), and we found exactly the same term U . In this case, we gave an argument that using a more precise cutoff on energy might cancel this term. The energy, unlike the worldsheet momentum, is a gauge-invariant physical quantity. This led us in section 4.2 to consider cutoffs on the Euclidean momentum of $|K| < \Lambda^2 - m^2$, with $m^2 = \frac{1}{4}$ or 1 for light or heavy modes. Then because of a cancellation of Λ^2 divergences, the final result changes by terms of the form $\gamma p_-^2 / \pi \times (\text{rational})$. It is difficult however to implement an independent cutoff on the heavy and light modes in the bubble diagram. We gave one rather Heath Robinson construction, involving two Pauli–Villars regulators, which manages to cancel half of the extra term U , see (41).

The reason that these regulators are needed at all is essentially that the leading interaction term \mathcal{L}_3 contains both left- and right-moving fields, and derivatives. This does not happen in the $AdS_5 \times S^5$ case, and as a result the analogous leading diagrams¹⁹ give only finite integrals.

We stress that these extra terms are not an artefact of the near-flat-space limit; the exact terms we find there are also obtained by taking the limit of our BMN results, as in (50). And further, the extra term U seen in the near-flat case scales like p_-^2 in the same way as the $\log 2$ term; thus, there is no obvious explanation for why they should not be seen in the giant magnon case [17], although the V terms would be invisible.

7.2. Group parameterization

Finally, it seems that these extra terms may be altered by the choice of the group parameterization used. A different parameterization was defined in [21], replacing our equation (5) with

$$G = \Lambda(t, \phi) e^{\mathbb{X}},$$

where \mathbb{X} contains both bosons and fermions. The resulting fields will be related in some complicated nonlinear way, but all *physical* results must agree. Using this parameterization,

¹⁸ This statement is strictly true in cases where the bubble alone is finite. However the argument that we can remove the $\log 2$ term holds for all the bosons (but fails for the light fermion).

¹⁹ By this we mean the two-loop ‘sunset’ diagram correcting the propagator in [7], since the leading interaction there is \mathcal{L}_4 .

\mathcal{L}_3 was calculated by the authors of [38], who were kind enough to share this term with us. This allows us to repeat the bubble calculation, and again we find an extra term. But the result is different: for the bubbles correcting the light boson, equivalent to (23) + (26), we now obtain

$$\delta m_{\text{bubble}}^2 = -\frac{1}{16\pi}\gamma p_-^2 + \frac{7}{128\pi}\gamma p_-^2 = \frac{-1}{128\pi}\gamma p_-^2. \quad (51)$$

This involves a cancellation of $1/\epsilon$ terms in the fermionic piece which is as nontrivial as those from our parameterization²⁰.

The quartic term \mathcal{L}_4 has not been calculated; thus, we cannot repeat the tadpole calculations. But we note that in the integrals used, (28) has no terms $\frac{1}{\pi}$, and thus the tadpoles never contribute to the terms of the form $\gamma p_-^2/\pi \times (\text{rational})$. So this result should survive in the complete answer.

We do not wish to place too much emphasis on (51), which depends on many details of the fermions. We can however make the simpler observation that the bosonic term in \mathcal{L}_3 is the same in both parameterizations. This term alone makes the bubble term for the $\langle yy \rangle$ correction nonzero, (44), producing an extra term not expected from the dispersion relation (43).

7.3. Interpretation of the heavy modes

The heavy modes are something of a puzzle in this example of the AdS/CFT correspondence. They are not present in the Bethe ansatz description, although each one can be made as a simple superposition of two light modes. But from the point of view of classical strings on $AdS_4 \times CP^3$, the heavy (bosonic) modes are simply four of the eight transverse spatial dimensions.

In [21] Zarembo began the investigation of what loop corrections can teach us about this puzzle. With the corrections that we find, we can make the simpler observation that the heavy mode becomes heavier than a pair of light modes, and is thus kinematically allowed to decay. We discussed this further in section 5.3.

7.4. Parallels to [4]

This recent paper has some similarities to ours. First of all note that the $1/R^2$ corrections computed there are analogous to the $1/\sqrt{\lambda}$ corrections computed here, since $1/\sqrt{\lambda} \sim \alpha'/R^2$. Both papers find a cubic interaction (which was absent in the $AdS_5 \times S^5$ case), and while in [4] this is at order $1/R$, it only contributes at $1/R^2$, much like our bubble diagram needing two cubic vertices.

Some of the results in both papers are small- p limits of the giant magnon case studied in [17]. In particular, both can see the $-\log 2/2\pi$ term seen there, and by altering the prescription can remove this (although here subject to the limitations mentioned above).

However, an important difference is that this paper is strictly in infinite volume, $J = \infty$, whereas [4] computes also some finite- J effects. (For instance we believe that equation (61) there is a Lüscher F-term.)

Acknowledgments

For discussions of this work, we thank Dmitri Bykov, Valentina Giangreco M Puletti, Tristan McLoughlin, Sameer Murthy, Jeff Murugan, Olof Ohlsson Sax and especially Konstantin Zarembo. We would also like to thank Olof and Tristan for comments on the manuscript. We

²⁰ For the BMN case of this we see terms that reduce to these, plus terms free of p_1 including $1/\epsilon$ terms, much like those in section 6.

are very grateful to the authors of [38], Chrysostomos Kalousios, Cristian Vergu and Anastasia Volovich, for sharing their cubic interaction term with us. PS would also like to thank Nordita for hospitality while working on this project. MCA also thanks, in TIFR tradition, the people of India for their support of basic science.

Appendix A. Fermionic matrices and notation

A given matrix M in $\mathfrak{osp}(2, 2|6)$ can be represented as a 10×10 super matrix as

$$M = \begin{pmatrix} X & \theta \\ \eta & Y \end{pmatrix},$$

where the bosonic blocks satisfy

$$X^t = -\mathbb{C}_4 X \mathbb{C}_4, \quad X^\dagger = -\Gamma_0 X \Gamma_0, \quad Y^t = -Y, \quad Y^* = Y,$$

and X corresponds to $\mathfrak{sp}(2, 2)$ and Y to $\mathfrak{so}(6)$. The odd blocks are fermionic, in the sense of having Grassmanian matrix elements, and satisfy

$$\eta = -\theta^t \mathbb{C}_4, \quad \theta^* = \mathbb{C}_4 \theta.$$

The matrices \mathbb{C}_4 and Γ_0 are charge conjugation and Γ -matrices of the AdS_4 space and are given explicitly in [13]. After kappa gauge fixing [26], the fermionic 4×6 block θ is given by²¹

$$\theta = \frac{1}{\sqrt{2}} \begin{pmatrix} \theta_{1,1} & \theta_{1,2} & \theta_{1,3} & \theta_{1,4} & i\theta_{1,4} & i\theta_{1,3} \\ \theta_{2,1} & \theta_{2,2} & \theta_{2,3} & \theta_{2,4} & i\theta_{2,4} & i\theta_{2,3} \\ \theta_{2,1}^\dagger & \theta_{2,2}^\dagger & \theta_{2,3}^\dagger & \theta_{2,4}^\dagger & -i\theta_{2,4}^\dagger & -i\theta_{2,3}^\dagger \\ -\theta_{1,1}^\dagger & -\theta_{1,2}^\dagger & -\theta_{1,3}^\dagger & -\theta_{1,4}^\dagger & i\theta_{1,4}^\dagger & i\theta_{1,3}^\dagger \end{pmatrix}.$$

The eight independent components are given by²²

$$\begin{aligned} \theta_{1,1} &= \frac{1}{\sqrt{2}}(\bar{\psi}_{-2} - \bar{\psi}_{+2} + \psi_-^1 + \psi_+^1), & \theta_{1,2} &= \frac{i}{\sqrt{2}}(\bar{\psi}_{-2} - \bar{\psi}_{+2} - \psi_-^1 - \psi_+^1), \\ \theta_{2,1} &= \frac{1}{\sqrt{2}}(-\bar{\psi}_{-1} + \bar{\psi}_{+1} + \psi_-^2 + \psi_+^2), & \theta_{2,2} &= \frac{i}{\sqrt{2}}(-\bar{\psi}_{-1} + \bar{\psi}_{+1} - \psi_-^2 - \psi_+^2), \\ \theta_{1,3} &= \frac{1}{2\sqrt{2}}(i(s_-)_3^1 - i(s_-)_4^1 + (s_+)_3^1 - (s_+)_4^1), \\ \theta_{1,4} &= \frac{1}{2\sqrt{2}}((s_-)_3^1 + (s_-)_4^1 - i(s_+)_3^1 - i(s_+)_4^1), \\ \theta_{2,3} &= \frac{1}{2\sqrt{2}}(i(s_-)_3^2 - i(s_-)_4^2 + (s_+)_3^2 - (s_+)_4^2), \\ \theta_{2,4} &= \frac{1}{2\sqrt{2}}((s_-)_3^2 + (s_-)_4^2 - i(s_+)_3^2 - i(s_+)_4^2). \end{aligned} \quad (\text{A.1})$$

In the Lagrangians (4) and (6), we single out the various graded parts of the current, $A = -G^{-1} dG$, using an automorphism Ω

$$\Omega(M) = \Upsilon M \Upsilon^{-1}, \quad \Omega(M^{(k)}) = i^k M,$$

²¹ Note that we have performed $\theta \rightarrow \Gamma_0 \theta$ and a simple rescaling with $1/\sqrt{2}$ compared to [13].

²² We will denote the AdS_4 $SU(2)$ with indices in the set $\{1, 2\}$ and the CP_3 $SU(2)$ with indices in the set $\{3, 4\}$. We also adopt the somewhat sloppy notation $\bar{\chi} = \chi^\dagger$.

where Υ is a constant matrix satisfying $\Upsilon^2 = -1$. For (6), the auxiliary field Π is given by

$$\Pi = \frac{i}{2}\Pi_+\Sigma_+ + \frac{i}{4}\Pi_-\Sigma_- + \Pi_t,$$

where

$$\Pi_t = \begin{pmatrix} \frac{i}{2}\Pi_i^{(z)}\Gamma_i & 0 \\ 0 & \frac{1}{2}\Pi^{(y)}T_5 + \Pi_i^{(\omega)}\tau_i + \bar{\Pi}_i^{(\bar{\omega})}\bar{\tau}_i \end{pmatrix}.$$

Once again, we point to [13] for the explicit representation of the various matrices. However, note that a nice property of Π is that it projects to $M^{(2)}$ for even elements,

$$\text{Str}\Pi A^{(2)} = \text{Str}\Pi A.$$

Appendix B. Simplifying the quartic Lagrangian

After performing the near-flat-space boost from the quartic BMN Lagrangian, the pure fermionic quartic piece comes out as

$$\begin{aligned} \mathcal{L}_{FF} = & -\frac{1}{16}((s_-)^\beta_a(s_-)^\delta_\beta(s_+)^\alpha_\gamma\partial_-(s_-)^\gamma_d - (s_-)^\beta_a(s_-)^\alpha_\gamma(s_+)^\delta_\beta\partial_-(s_-)^\gamma_d) - \frac{1}{8}(\bar{\psi}_-\psi_-)^2 \\ & + \frac{1}{8}(\bar{\psi}_-\overset{\leftrightarrow}{\partial}_-\psi_-)(\bar{\psi}_-\overset{\leftrightarrow}{\partial}_+\psi_-) \\ & + \frac{1}{4}(\bar{\psi}_-\partial_+\psi_-\bar{\psi}_-\partial_-\psi_- + \partial_+\bar{\psi}_-\psi_-\partial_-\bar{\psi}_-\psi_- \\ & - \bar{\psi}_-\psi_-\partial_-\bar{\psi}_-\partial_+\psi_- - \partial_+\bar{\psi}_-\partial_-\psi_-\bar{\psi}_-\psi_-) \\ & + \frac{i}{16}\bar{\psi}_-\overset{\leftrightarrow}{\partial}_-\psi_-(\bar{\psi}_+\psi_+ + \bar{\psi}_-\psi_+) - \frac{i}{16}\bar{\psi}_-\psi_-\partial_-(\bar{\psi}_+\psi_+ - \bar{\psi}_-\psi_+) \\ & + \frac{i}{2}\bar{\psi}_-\psi_-(\bar{\psi}_+\partial_-\psi_- - \partial_-\bar{\psi}_-\psi_+) + \frac{3i}{8}(\bar{\psi}_-\partial_-\psi_-\bar{\psi}_+\psi_- - \partial_-\bar{\psi}_-\psi_-\bar{\psi}_-\psi_+) \\ & + \frac{1}{8}\left[-\partial_+(s_-)^\alpha_a(s_-)^\alpha_c\bar{\psi}_-{}_a\partial_-\psi_-^c + \frac{1}{2}\partial_+(s_-)^\alpha_a(s_-)^\alpha_c\partial_-\bar{\psi}_-{}_a\psi_-^c \right. \\ & + \frac{1}{2}\partial_+(s_-)^\alpha_a\partial_-(s_-)^\alpha_c\bar{\psi}_-{}_a\psi_-^c + \frac{1}{2}(s_-)^\alpha_a\partial_+(s_-)^\alpha_c\bar{\psi}_-{}_a\partial_-\psi_-^c \\ & - (s_-)^\alpha_a\partial_+(s_-)^\alpha_c\partial_-\bar{\psi}_-{}_a\psi_-^c + \frac{1}{2}\partial_-(s_-)^\alpha_a\partial_+(s_-)^\alpha_c\bar{\psi}_-{}_a\psi_-^c \\ & - \frac{1}{4}(s_-)^\alpha_a(s_-)^\alpha_c\bar{\psi}_-{}_a\psi_-^c + \frac{1}{2}(s_-)^\alpha_a(s_-)^\alpha_c\partial_+\bar{\psi}_-{}_a\partial_-\psi_-^c \\ & + \frac{1}{2}(s_-)^\alpha_a(s_-)^\alpha_c\partial_-\bar{\psi}_-{}_a\partial_+\psi_-^c - \frac{1}{2}(s_-)^\alpha_a\partial_-(s_-)^\alpha_c\bar{\psi}_-{}_a\partial_+\psi_-^c \\ & - \frac{1}{2}\partial_-(s_-)^\alpha_a(s_-)^\alpha_c\partial_+\bar{\psi}_-{}_a\psi_-^c - 2(s_-)^\alpha_a(s_+)^\alpha_c\partial_-\bar{\psi}_-{}_a\psi_-^c \\ & - 2(s_+)^\alpha_a(s_-)^\alpha_c\bar{\psi}_-{}_a\partial_-\psi_-^c + i\partial_-(s_-)^\alpha_a(s_-)^\alpha_c\bar{\psi}_+{}_a\psi_-^c \\ & \left. - i(s_-)^\alpha_a\partial_-(s_-)^\alpha_c\bar{\psi}_-{}_a\psi_+^c\right]. \end{aligned} \quad (\text{B.1})$$

The first thing to note is that, using total derivatives, the piece quartic in ψ and s and involving one left mover can be written as

$$\begin{aligned} & -\frac{1}{24}(s_-)^\beta_a(s_-)^\delta_\beta(s_-)^\gamma_d\partial_-(s_+)^\alpha_\gamma + \frac{i}{8}(3\bar{\psi}_-\partial_-\psi_- + \partial_-\bar{\psi}_-\psi_-)\bar{\psi}_-\psi_+ \\ & -\frac{i}{8}(3\partial_-\bar{\psi}_-\psi_- + \bar{\psi}_-\partial_-\psi_-)\bar{\psi}_+\psi_- - \frac{i}{2}\bar{\psi}_-\psi_-(\partial_-\bar{\psi}_+\psi_- - \bar{\psi}_-\partial_-\psi_+). \end{aligned} \quad (\text{B.2})$$

If we now perform the following redefinitions of the fermions (and corresponding ones for conjugated ψ_{\pm}):

$$\begin{aligned}\delta\psi_+^a &= -\frac{1}{2}\psi_-^a\bar{\psi}_-\psi_-, & \delta\psi_-^a &= -\frac{i}{4}\psi_-^a(3\partial_-\bar{\psi}_-\psi_- + \bar{\psi}_-\partial_-\psi_-), \\ \delta(s_+)_a^\gamma &= -\frac{i}{24}(s_-)_a^\beta(s_-)_\beta^d(s_-)_d^\gamma, & & \end{aligned}$$

then the mass term in the quadratic Lagrangian (8) removes (B.2). However, the kinetic term induces additional terms giving

$$\begin{aligned}\mathcal{L}_{FF} &= \frac{3}{8}(\bar{\psi}_-\psi_-)^2 - \frac{1}{4}\bar{\psi}_-\psi_-(\partial_+\bar{\psi}_-\partial_-\psi_- + \partial_-\bar{\psi}_-\partial_+\psi_-) \\ &\quad - \frac{3}{8}(\bar{\psi}_-\partial_+\psi_-\bar{\psi}_-\partial_-\psi_- + \partial_+\bar{\psi}_-\psi_-\partial_-\bar{\psi}_-\psi_- \\ &\quad + \partial_-\bar{\psi}_-\psi_-\bar{\psi}_-\partial_+\psi_- + \partial_+\bar{\psi}_-\psi_-\bar{\psi}_-\partial_-\psi_-) \\ &\quad - \frac{1}{24}(s_-)_a^\beta(s_-)_\beta^d(s_-)_d^\gamma(s_-)_\gamma^a + \dots, \end{aligned}$$

where the dots denote the unchanged terms mixing s_{\pm} and ψ_{\pm} . The mixing terms linear in left movers

$$\begin{aligned}\frac{1}{4}\left((s_+)_c^\alpha(s_-)_\alpha^a\partial_-\bar{\psi}_{-a}\psi_-^c - (s_+)_\alpha^a(s_-)_c^\alpha\bar{\psi}_{-a}\partial_-\psi_-^c\right. \\ \left.+ \frac{i}{2}\bar{\psi}_{+a}\partial_-(s_-)_\alpha^a(s_-)_c^\alpha\psi_-^c - \frac{i}{2}(s_-)_\alpha^a\partial_-(s_-)_c^\alpha\bar{\psi}_{-a}\psi_+^c\right) \end{aligned} \quad (\text{B.3})$$

can be removed by performing a similar field redefinition. That is, we introduce

$$\delta(s_-)_\alpha^c = -\frac{i}{4}(s_-)_\alpha^a(\partial_-\bar{\psi}_{-a}\psi_-^c + \bar{\psi}_{-a}^c\partial_-\psi_{-a}), \quad \delta\psi_-^a = \frac{i}{4}\partial_-(s_-)_\alpha^a(s_-)_c^\alpha\psi_-^c$$

which removes (B.3) at the cost of additional quartic terms. Adding it all together gives the \mathcal{L}_{FF} presented in (20).

Appendix C. Expansions in dimensional regularization

In this appendix, we consider some changes to our calculation which would produce the terms of the form $\gamma p_-^2/\pi \times (\text{rational})$, thus altering the terms U in (50). While we believe that what we have done is correct, we include these other options as a matter of curiosity.

The integrals we write in (25) for $s = 1, 2$ are closely related to these two:²³

$$\begin{aligned}\int \frac{d^d k}{(2\pi)^d} \frac{k_\mu k_\nu}{[k^2 - \Delta]^n} &= \frac{-i}{(4\pi)^{d/2}} \frac{g_{\mu\nu}}{2} \left(\frac{1}{\Delta}\right)^{n-1-d/2} \frac{\Gamma(n-1-\frac{d}{2})}{\Gamma(n)}, \\ \int \frac{d^d k}{(2\pi)^d} \frac{k_\mu k_\nu k_\xi k_\varpi}{[k^2 - \Delta]^n} &= \frac{i}{(4\pi)^{d/2}} \frac{g_{\mu\nu}g_{\xi\varpi} + g_{\mu\xi}g_{\nu\varpi} + g_{\mu\varpi}g_{\nu\xi}}{4} \left(\frac{1}{\Delta}\right)^{1-d/2} \frac{\Gamma(n-2-\frac{d}{2})}{\Gamma(n)}. \end{aligned} \quad (\text{C.1})$$

Tracing produces the factors of $d/2$ and $\frac{d}{4}(d+2)$, which in (25) are written as a quotient of gamma functions:

$$\begin{aligned}g_{\mu\nu} \frac{g_{\mu\nu}}{2} &= \frac{d}{2} = \frac{\Gamma(\frac{d}{2}+1)}{\Gamma(\frac{d}{2})} \\ g_{\mu\nu}g_{\xi\varpi} \frac{g_{\mu\nu}g_{\xi\varpi} + g_{\mu\xi}g_{\nu\varpi} + g_{\mu\varpi}g_{\nu\xi}}{4} &= \frac{1}{4}d(d+2) = \frac{\Gamma(\frac{d}{2}+2)}{\Gamma(\frac{d}{2})}. \end{aligned} \quad (\text{C.2})$$

²³ Here we note that [39] has a misprint in (A.47), which is the $s = 2$ case of our (25), as well as in (A.51).

These factors are multiplied by the divergent gamma function $\Gamma(\frac{\epsilon}{2})$ or $\Gamma(-1 + \frac{\epsilon}{2})$, and to get the finite terms shown in (25) we have expanded these (and the other non-divergent factors) up to order ϵ . This seems like the most natural and correct thing to do, when starting from integrals with k^2 or $(k^2)^2$ in the numerator. There are however some other options.

2. We could expand neither of the factors in (C.2). In other words, use both of the integrals (C.1) rather than (25), and trace them only at the end. This produces changes to the various $1/\pi$ terms, which (in the BMN case) lead to the following:

$$\begin{aligned}\delta m^2 &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{13}{12\pi} p_1^2 + \frac{19}{64\pi} - \frac{9a}{32\pi} \right) \quad \text{for } \omega \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(+\frac{17}{96\pi} p_1^1 - \frac{35}{768\pi} - \frac{9a}{32\pi} \right) \psi \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{1}{4\pi} p_1^2 - \frac{17}{48\pi} - \frac{i}{4} \right) y \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{11}{12\pi} p_1^2 + \frac{1}{12\pi} - \frac{3a}{4\pi} \right) z.\end{aligned}$$

None of these are preferable to our results above. Note the appearance of the parameter a of the generalized light-cone gauge in some of these, although only in the constant term (thus not visible in the near-flat limit).

3. We could expand the $(d+2)$ but not the $d/2$ in (C.2). This would mean in some sense that we are keeping the same indices $\mu\nu$ outside the integral in all cases—we use the first integral in (C.1) but trace the second integral once, like

$$\begin{aligned}g_{\mu\nu} g_{\xi\varpi} \frac{g_{\mu\nu} g_{\xi\varpi} + g_{\mu\xi} g_{\nu\varpi} + g_{\mu\varpi} g_{\nu\xi}}{4} &= \frac{g_{\mu\nu}(d+2)}{4} \\ \Rightarrow \int \frac{d^d k}{(2\pi)^d} \frac{k_\mu k_\nu k^2}{[k^2 - \Delta]^n} &= \frac{i}{(4\pi)^{d/2}} \frac{g_{\mu\nu}(d+2)}{4} \left(\frac{1}{\Delta} \right)^{1-d/2} \frac{\Gamma(n-2-\frac{d}{2})}{\Gamma(n)}.\end{aligned}$$

This third option leads to the following results:

$$\begin{aligned}\delta m^2 &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{1}{2\pi} p_1^2 + \frac{5}{64\pi} - \frac{9a}{32\pi} \right) \quad \text{for } \omega \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(\frac{3}{8\pi} p_1^1 - \frac{1}{384\pi} - \frac{9a}{32\pi} \right) \psi \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{1}{4\pi} p_1^2 - \frac{17}{48\pi} - \frac{i}{4} \right) y \\ &= -\frac{\log 2}{\pi} \gamma p_1^2 + \gamma \left(-\frac{3}{4\pi} p_1^2 - \frac{5}{24\pi} - \frac{3a}{4\pi} \right) z.\end{aligned}$$

Again these are not any better than our initial results.

References

- [1] Maldacena J M and Swanson I 2007 Connecting giant magnons to the pp-wave: An interpolating limit of $AdS_5 \times S^5$ *Phys. Rev. D* **76** 026002
- [2] McLoughlin T, Roiban R and Tseytlin A A 2008 Quantum spinning strings in $AdS_4 \times CP^3$: testing the Bethe ansatz proposal *J. High Energy Phys.* **JHEP11(2008)069**
- [3] Gromov N and Mikhaylov V 2009 Comment on the scaling function in $AdS_4 \times CP^3$ *J. High Energy Phys.* **JHEP04(2009)083**

- [4] Astolfi D, Puletti V G M, Grignani G, Harmark T and Orselli M 2011 Finite-size corrections for quantum strings on $AdS_4 \times CP^3$ arXiv:1101.0004v2 [hep-th]
- [5] Berenstein D E, Maldacena J M and Nastase H S 2002 Strings in flat space and pp waves from $\mathcal{N} = 4$ super yang mills *J. High Energy Phys.* **JHEP04(2002)013**
- [6] Klose T and Zarembo K 2007 Reduced sigma-model on $AdS_5 \times S^5$: one-loop scattering amplitudes *J. High Energy Phys.* **JHEP02(2007)071**
- [7] Klose T, McLoughlin T, Minahan J A and Zarembo K 2007 World-sheet scattering in $AdS_5 \times S^5$ at two loops *J. High Energy Phys.* **JHEP08(2007)051**
- [8] Ahn C and Nepomechie R I 2010 Review of AdS/CFT integrability, chapter iii.2: Exact world-sheet S-matrix arXiv:1012.3991v3 [hep-th]
- [9] Hofman D M and Maldacena J M 2006 Giant magnons *J. Phys. A: Math. Gen.* **39** 13095–118
- [10] Aharony O, Bergman O, Jafferis D L and Maldacena J M 2008 $\mathcal{N} = 6$ superconformal Chern–Simons-matter theories, M2-branes and their gravity duals *J. High Energy Phys.* **JHEP10(2008)091**
- [11] Klose T 2010 Review of AdS/CFT integrability, chapter iv.3: $\mathcal{N} = 6$ Chern–Simons and strings on $AdS_4 \times CP^3$ arXiv:1012.3999v3 [hep-th]
- [12] Kreuzer M, Rashkov R C and Schimpf M 2009 Near flat space limit of strings on $AdS_4 \times CP^3$ *Eur. Phys. J. C* **60** 471–80
- [13] Sundin P 2010 On the worldsheet theory of the type IIA $AdS_4 \times CP^3$ superstring *J. High Energy Phys.* **JHEP04(2010)014**
- [14] McLoughlin T and Roiban R 2008 Spinning strings at one-loop in $AdS_4 \times CP^3$ *J. High Energy Phys.* **JHEP12(2008)101**
- [15] Alday L F, Arutyunov G and Bykov D 2008 Semiclassical quantization of spinning strings in $AdS_4 \times CP^3$ *J. High Energy Phys.* **JHEP11(2008)089**
- [16] Krishnan C 2008 AdS_4/CFT_3 at one loop *J. High Energy Phys.* **JHEP09(2008)092**
- [17] Abbott M C, Aniceto I and Bombardelli D 2010 Quantum strings and the AdS_4/CFT_3 interpolating function *J. High Energy Phys.* **JHEP12(2010)040**
- [18] Shenderovich I 2008 Giant magnons in AdS_4/CFT_3 : dispersion, quantization and finite-size corrections arXiv:0807.2861v3 [hep-th]
- [19] Astolfi D, Puletti V G M, Grignani G, Harmark T and Orselli M 2008 Finite-size corrections in the $su(2) \times su(2)$ sector of type IIA string theory on $AdS_4 \times CP^3$ *Nucl. Phys. B* **810** 150–73
- [20] Sundin P 2009 The $AdS_4 \times CP^3$ string and its Bethe equations in the near plane wave limit *J. High Energy Phys.* **JHEP02(2009)046**
- [21] Zarembo K 2009 Worldsheet spectrum in AdS_4/CFT_3 correspondence *J. High Energy Phys.* **JHEP04(2009)135**
- [22] Arutyunov G and Frolov S 2008 Superstrings on $AdS_4 \times CP^3$ as a coset sigma-model *J. High Energy Phys.* **JHEP09(2008)129**
- [23] Stefanski B 2009 Green–Schwarz action for type IIA strings on $AdS_4 \times CP^3$ *Nucl. Phys. B* **808** 80–7
- [24] Arutyunov G, Frolov S and Zamaklar M 2007 Finite-size effects from giant magnons *Nucl. Phys. B* **778** 1–35
- [25] Klose T, McLoughlin T, Roiban R and Zarembo K 2007 Worldsheet scattering in $AdS_5 \times S^5$ *J. High Energy Phys.* **JHEP03(2007)094**
- [26] Bykov D 2010 Symmetry algebra of the $AdS_4 \times CP^3$ superstring *Teor. Mat. Fiz.* **163** 496–510
- [27] Astolfi D, Puletti V G M, Grignani G, Harmark T and Orselli M 2010 Full Lagrangian and Hamiltonian for quantum strings on $AdS_4 \times CP^3$ in a near plane wave limit *J. High Energy Phys.* **JHEP04(2010)079**
- [28] Sundin P 2010 Perturbative quantization of superstring theory in anti de Sitter spaces *PhD Thesis* Humboldt University
- [29] Frolov S, Plefka J and Zamaklar M 2006 The $AdS_5 \times S^5$ superstring in light-cone gauge and its Bethe equations *J. Phys. A: Math. Gen.* **39** 13037–82
- [30] Arutyunov G and Frolov S 2009 Foundations of the $AdS_5 \times S^5$ superstring: part I *J. Phys. A: Math. Theor.* **42** 254003
- [31] Leibbrandt G 1975 Introduction to the technique of dimensional regularization *Rev. Mod. Phys.* **47** 849–76
- [32] Pauli W and Villars F 1949 On the invariant regularization in relativistic quantum theory *Rev. Mod. Phys.* **21** 434–44
- [33] Bandres M A and Lipstein A E 2010 One-loop corrections to type IIA string theory in $AdS_4 \times CP^3$ *J. High Energy Phys.* **JHEP04(2010)059**
- [34] Hollowood T J and Miramontes J L 2009 A new and elementary CP^n dyonic magnon *J. High Energy Phys.* **JHEP08(2009)109**
- [35] Abbott M C, Aniceto I and Sax O O 2009 Dyonic giant magnons in CP^3 : strings and curves at finite *J. Phys. Rev. D* **80** 026005

- [36] Gromov N and Vieira P 2008 The AdS_4/CFT_3 algebraic curve *J. High Energy Phys.* [JHEP02\(2008\)040](#)
- [37] Bellucci S 1987 Finite renormalizations of $OSP(n, 4)$ supersymmetry *Z. Phys. C* **34** 249–54
- [38] Kalousios C, Vergu C and Volovich A 2009 Factorized tree-level scattering in $AdS_4 \times CP^3$ *J. High Energy Phys.* [JHEP09\(2009\)049](#)
- [39] Peskin M E and Schroeder D V 1995 *An Introduction to Quantum Field Theory* (Reading, MA: Addison-Wesley)