

# Lectures on hydrodynamics - Part III: Causal dissipative hydrodynamics

**Denes Molnar**

RIKEN/BNL Research Center & Purdue University

**Goa Summer School**


September 8-12, 2008, **International Centre**, Dona Paula, Goa, India

# Outline

- Why **dissipative**
- **Viscosities in QCD**
- **Dissipative hydro EOMs - Navier-Stokes, Israel-Stewart theory**
- **What are the right equations? - cross-check from covariant transport**
- **Current state of art and open problems**

# Hydrodynamics

- describes a system near local equilibrium
- long-wavelength, long-timescale dynamics, driven by conservation laws
- in heavy-ion physics: mainly used for the plasma stage of the collision




nontrivial how hydrodynamics can be applicable at such microscopic scales

# Shear viscosity

1687 - I. Newton (Principia)

$$T_{xy} = -\eta \frac{\partial u_x}{\partial y}$$

acts to reduce velocity gradients



1985 - quantum mechanics:  $\Delta E \cdot \Delta t \geq \hbar/2$

+ kinetic theory:  $T \cdot \lambda_{MFP} \geq \hbar/3$  Gyulassy & Danielewicz, PRD 31 ('85)


$$\eta \approx 4/5 \cdot T/\sigma_{tr} , \quad \text{entropy } s \approx 4n$$

gives minimal viscosity:  $\eta/s = \frac{\lambda_{tr} T}{5} \geq \hbar/15$

2004 - string theory AdS/CFT:  $\eta/s \geq \hbar/4\pi$  Policastro, Son, Starinets, PRL87 ('02)  
Kovtun, Son, Starinets, PRL94 ('05)

revised to  $4\hbar/(25\pi)$  Brigante et al, arXiv:0802.3318

$\sim 1/(4\pi)$  bound **conjectured** universal - at least no other known substance comes within a factor **10** Kovtun, Son, Starinets, PRL94 ('05):



**(perhaps strongly-interacting cold atoms...)**


# Shear viscosity in QCD

$$\eta = \lim_{\omega \rightarrow 0} \frac{1}{2\omega} \int dt d^3x e^{i\omega t} \langle [T_{xy}(x), T_{xy}(0)] \rangle$$

**perturbative QCD:**  $\eta/s \sim 1$ ,      **lattice QCD:** correlator very noisy

Nakamura & Sakai, NPA774, 775 ('06):

Meyer, PRD76, 101701 ('07)



**upper bounds:**

$$\eta/s(T=1.65T_c) < 0.96$$

$$\eta/s(T=1.24T_c) < 1.08$$

**best estimate:**

$$\eta/s(T=1.65T_c) < 0.13 \pm 0.03$$

$$\eta/s(T=1.24T_c) < 0.10 \pm 0.05$$


**many practitioners regard these VERY preliminary**

# Bulk viscosity in QCD

$$\zeta = \lim_{\omega \rightarrow 0} \frac{1}{18\omega} \int dt d^3x e^{i\omega t} \langle [T_\mu^\mu(\vec{x}, t), T_\mu^\mu(0)] \rangle$$

**perturbative QCD:**  $\zeta/s \sim 0.02\alpha_s^2$  is tiny Arnold, Dogan, Moore, PRD74 ('06)

**from  $\varepsilon - 3p > 0$ :** Kharzeev & Tuchin, arXiv:0705.4280v2



**on lattice:** Meyer, arXiv:0710.3717

**best estimates:**

$$\eta/s(T=1.65T_c) \sim 0 - 0.015$$

$$\eta/s(T=1.24T_c) \sim 0.06 - 0.1$$

$$\eta/s(T=1.02T_c) \sim 0.2 - 2.7$$

**many practitioners regard these as well VERY preliminary**

If we can quantify dissipative effects on heavy ion observables, we could constrain the viscosities. But cannot use ideal hydro, which has no dissipation.

## Two ways to study dissipative effects in heavy-ion collisions

### - causal dissipative hydrodynamics

Israel, Stewart; ... Muronga, Rischke; Teaney et al; Romatschke et al; Heinz et al, DM & Huovinen

**flexible in macroscopic properties**

**numerically cheaper**

### - covariant transport

Israel, de Groot,... Zhang, Gyulassy, DM, Pratt, Xu, Greiner...

**completely causal and stable**

**fully nonequilibrium → interpolation to break-up stage**

## Several active groups

- Paul Romatschke et al
- Huichao Song & Ulrich Heinz
- Derek Teaney & Kevin Dusling
- DM & Pasi Huovinen
- Takeshi Kodama & Tomo Koide et al
- ...

no public codes (yet)

State of the art is 2+1D calculations (with Bjorken boost invariance)

# Relativistic dissipative hydro

Decompose energy-momentum tensor and currents using a flow field  $u^\mu(x)$

In local rest frame (LR) (where  $u_{LR}^\mu = (1, \vec{0})$ ),

$$T_{LR}^{\mu\nu} = \begin{pmatrix} \varepsilon & h_x & h_y & h_z \\ h_x & p + \pi_{xx} + \Pi & \pi_{xy} & \pi_{xz} \\ h_y & \pi_{xy} & p + \pi_{yy} + \Pi & \pi_{yz} \\ h_z & \pi_{xz} & \pi_{yz} & p + \pi_{zz} + \Pi \end{pmatrix}, \quad N_{LR}^\mu = (n, \vec{V})$$

$\vec{h}(x)$  - energy flow,  $\Pi(x)$  - bulk pressure,  $\pi_{ij}(x)$  - shear stress

In general frame

$$T^{\mu\nu} = (\varepsilon + p + \Pi)u^\mu u^\nu - (p + \Pi)g^{\mu\nu} + (u^\mu h^\nu + u^\nu h^\mu) + \pi^{\mu\nu}$$

$$N^\mu = nu^\mu + V^\mu \quad (u^\mu h_\mu = 0 = u^\mu V_\mu, \quad u^\mu \pi_{\mu\nu} = 0 = \pi_{\mu\nu} u^\nu, \quad \pi_\nu^\nu = 0)$$

So far  $u^\mu(x)$  is arbitrary. Most common choices:

- **Eckart**: no particle flow in LR  $\rightarrow \vec{V} = 0$ ,  $u^\mu = N^\mu / \sqrt{N_\alpha N^\alpha}$
- **Landau**: no energy flow in LR  $\rightarrow \vec{h} = 0$ ,  $u^\mu = u^\nu T^{\mu\nu} / \sqrt{u_\alpha T^{\alpha\beta} T_{\beta\gamma} u^\gamma}$

# Dissipative hydrodynamics

**relativistic Navier-Stokes hydro: small corrections linear in gradients** [Landau]

$$T_{NS}^{\mu\nu} = T_{ideal}^{\mu\nu} + \eta(\nabla^\mu u^\nu + \nabla^\nu u^\mu - \frac{2}{3}\Delta^{\mu\nu}\partial^\alpha u_\alpha) + \zeta\Delta^{\mu\nu}\partial^\alpha u_\alpha$$
$$N_{NS}^\nu = N_{ideal}^\nu + \kappa \left(\frac{n}{\varepsilon + p}\right)^2 \nabla^\nu \left(\frac{\mu}{T}\right)$$

where  $\Delta^{\mu\nu} \equiv u^\mu u^\nu - g^{\mu\nu}$ ,  $\nabla^\mu = \Delta^{\mu\nu}\partial_\nu$

$\eta, \zeta$  shear and bulk viscosities,  $\kappa$  heat conductivity

**Equation of motion:**  $\partial_\mu T^{\mu\nu} = 0$ ,  $\partial_\mu N^\mu = 0$

two problems:

**parabolic equations**  $\rightarrow$  **acausal** Müller ('76), Israel & Stewart ('79) ...

**instabilities** Hiscock & Lindblom, PRD31, 725 (1985) ...

**As an illustration, consider heat flow in a static, incompressible fluid (Fourier)**

$$\partial_t T = \kappa \Delta T$$

**parabolic eqns.**

**Greens function is acausal (allows  $\Delta x > \Delta t$ )**

$$G(\vec{x}, t; \vec{x}_0, t_0) = \frac{1}{[4\pi\kappa(t - t_0)]^{3/2}} \exp\left[-\frac{(\vec{x} - \vec{x}_0)^2}{4\kappa(t - t_0)}\right]$$

—

**Adding a second-order time derivative makes it hyperbolic**

$$\tau \partial_t^2 T + \partial_t T = \kappa \Delta T$$

**Note, this is equivalent to a relaxing heat current**

$$\partial_\tau T = \vec{\nabla} \cdot \vec{j}, \quad \partial_t \vec{j} = -\frac{\vec{j} - \kappa \vec{\nabla} T}{\tau}$$

**The wave dispersion relation is  $\omega^2 + i\omega/\tau = \kappa k^2/\tau$ , i.e., now signals propagate at speeds  $c_s = \sqrt{\kappa/\tau}$  (at low frequencies), causal for large enough  $\tau$ .**

# Causal dissipative hydro

Bulk pressure  $\Pi$ , shear stress  $\pi^{\mu\nu}$ , heat flow  $q^\mu$  are dynamical quantities

$$T^{\mu\nu} \equiv T_{ideal}^{\mu\nu} + \pi^{\mu\nu} - \Pi\Delta^{\mu\nu}, \quad N^\mu \equiv N_{ideal}^\mu - \frac{n}{e+p}q^\mu$$

**Israel-Stewart: truncate entropy current at quadratic order** [Ann.Phys 100 & 118]

$$S^\mu = u^\mu \left[ s - \frac{1}{2T} (\beta_0\Pi^2 - \beta_1q_\nu q^\nu + \beta_2\pi_{\nu\lambda}\pi^{\nu\lambda}) \right] + \frac{q^\mu}{T} \left( \frac{\mu n}{\varepsilon + p} + \alpha_0\Pi \right) - \frac{\alpha_1}{T}\pi^{\mu\nu}q_\nu$$

in Landau frame. Note,  $\alpha_0 = \alpha_1 = \beta_0 = \beta_1 = \beta_2 = 0$  gives Navier-Stokes.

Impose  $\partial_\mu S^\mu \geq 0$  via a quadratic ansatz

$$T\partial_\mu S^\mu = \frac{\Pi^2}{\zeta} - \frac{q_\mu q^\mu}{\kappa_q T} + \frac{\pi_{\mu\nu}\pi^{\mu\nu}}{2\eta_s} \geq 0$$

E.g.,

$$T\partial_\mu S^\mu = \Pi X - q^\mu X_\mu + \pi^{\mu\nu} X_{\mu\nu}$$

will lead to equations of motion

$$\Pi = \zeta X, \quad q^\mu = \kappa T \Delta^{\mu\nu} X_\nu, \quad \pi^{\mu\nu} = 2\eta_s X^{\langle\mu\nu\rangle}$$

The resulting equations **relax** dissipative quantities on time scales

$$\tau_{\Pi}(e, n) = \beta_0 \zeta, \quad \tau_{\pi}(e, n) = 2\beta_2 \eta, \quad \tau_q(e, n) = \beta_1 \kappa T$$

toward values set by gradients - because not only first but also certain second derivatives are kept.

schematically

$$\dot{X} = -\frac{X - X_0}{\tau_X} + X_c$$

**restores causality (for not too small  $\tau_X$ )** telegraph eqn

**Splitting  $q_{\Pi}$  and  $q_{\pi}$  terms between heat and bulk, and heat and shear equations is ambiguous, and requires additional matter parameters  $a_0(\varepsilon, n)$ ,  $a_1(\varepsilon, n)$  to specify** Israel, Stewart... Huovinen & DM, arXiv:0808.0953

**Moreover, further terms that produce no entropy can be added, which are missed by the Israel-Stewart procedure.**

# Complete set of Israel-Stewart equations of motion

$$D\Pi = -\frac{1}{\tau_\Pi} (\Pi + \zeta \nabla_\mu u^\mu) \quad (1)$$

$$Dq^\mu = -\frac{1}{\tau_q} \left[ q^\mu + \kappa_q \frac{T^2 n}{\varepsilon + p} \nabla^\mu \left( \frac{\mu}{T} \right) \right] - u^\mu q_\nu D u^\nu \quad (2)$$

$$D\pi^{\mu\nu} = -\frac{1}{\tau_\pi} \left( \pi^{\mu\nu} - 2\eta \nabla^{\langle\mu} u^{\nu\rangle} \right) - (\pi^{\lambda\mu} u^\nu + \pi^{\lambda\nu} u^\mu) D u_\lambda \quad (3)$$

$$-\frac{1}{2} q^\mu \left( \nabla_\lambda u^\lambda + D \ln \frac{\beta_1}{T} \right) - \omega^{\mu\lambda} q_\lambda$$


$$-\frac{\alpha_0}{\beta_1} \nabla^\mu \Pi + \frac{\alpha_1}{\beta_1} (\partial_\lambda \pi^{\lambda\mu} + u^\mu \pi^{\lambda\nu} \partial_\lambda u_\nu) + \frac{a_0}{\beta_1} \Pi D u^\mu - \frac{a_1}{\beta_1} \pi^{\lambda\mu} D u_\lambda$$

$$-\frac{1}{2} \pi^{\mu\nu} \left( \nabla_\lambda u^\lambda + D \ln \frac{\beta_2}{T} \right) - 2\pi_\lambda^{\langle\mu} \omega^{\nu\rangle\lambda}$$


$$-\frac{\alpha_1}{\beta_2} \nabla^{\langle\mu} q^{\nu\rangle} + \frac{a'_1}{\beta_2} q^{\langle\mu} D u^{\nu\rangle} .$$

where  $A^{\langle\mu\nu\rangle} \equiv \frac{1}{2} \Delta^{\mu\alpha} \Delta^{\nu\beta} (A_{\alpha\beta} + A_{\beta\alpha}) - \frac{1}{3} \Delta^{\mu\nu} \Delta_{\alpha\beta} A^{\alpha\beta}$ ,  $\omega^{\mu\nu} \equiv \frac{1}{2} \Delta^{\mu\alpha} \Delta^{\nu\beta} (\partial_\beta u_\alpha - \partial_\alpha u_\beta)$

# Recent viscous hydro calculations disagreed



Romatschke & Romatschke, arxiv:0706.1522



Song & Heinz, arxiv:0709.0742

for  $\eta/s = 1/(4\pi)$ , **~20%** or **50+% elliptic flow reduction??**

Origin of difference is in WHICH TERMS are kept in Israel-Stewart eqns:

$$\begin{aligned} \dot{\pi}^{\mu\nu} &= -\frac{1}{\tau_\pi}(\pi^{\mu\nu} - 2\eta\nabla^{\langle\mu}u^{\nu\rangle}) - (u^\mu\pi^{\nu\alpha} + u^\nu\pi^{\alpha\mu})u_\alpha \\ &- \frac{1}{2}\pi^{\mu\nu}D_\alpha u^\alpha - \frac{1}{2}\pi^{\mu\nu}[\ln \frac{\beta_2}{T}] + 2\pi_\lambda^{\langle\mu}\omega^{\nu\rangle\lambda} \end{aligned} \quad (4)$$

Heinz et al neglected **terms in green**.

Which terms to keep?? Can only tell via comparing to a nonequilibrium theory.

# IS hydro and covariant transport

Israel-Stewart hydro can be derived from covariant transport through **Grad's 14-moment approximation**

$$f(x, p) \approx [1 + \tilde{C}_\alpha p^\alpha + C_{\alpha\beta} p^\alpha p^\beta] f_{eq}(x, p)$$

via taking the “1”,  $p^\nu$ , and  $p^\nu p^\alpha$  moments of the transport equation.

However, whereas Navier-Stokes came from a rigorous expansion in small deviations near local equilibrium retaining all powers of momentum (recall integral eqn from Part II), **the quadratic truncation in Grad's approach has no small control parameter.**

If relaxation effects important, NS and IS are different

**⇒ control against a nonequilibrium theory is crucial**

# Applicability of IS hydro

**Important to realize - in heavy ion physics applications, gradients  $\partial^\mu u^\nu/T$ ,  $|\partial^\mu e|/(Te)$ ,  $|\partial^\mu n|/(Tn)$  at early times  $\tau \sim 1$  fm are large  $\sim \mathcal{O}(1)$ , and therefore cannot be ignored.**

**Hydrodynamics may still apply, if viscosities are unusually small  $\eta/s \sim 0.1$ ,  $\zeta/s \sim 0.1$ , where  $s$  is the entropy density in local equilibrium. In that case, pressure corrections from Navier-Stokes theory still moderate**

$$\frac{\delta T_{NS}^{\mu\nu}}{p} \approx \left( 2 \frac{\eta_s}{s} \frac{\nabla^{\langle\mu} u^{\nu\rangle}}{T} + \frac{\zeta}{s} \frac{\nabla_\alpha u^\alpha}{T} \right) \frac{\varepsilon + p}{p} \sim \mathcal{O} \left( \frac{8\eta_s}{s}, \frac{4\zeta}{s} \right) . \quad (5)$$

**Heat flow effects can also be estimated based on**

$$\frac{\delta N_{NS}^\mu}{n} \approx \frac{\kappa_q T n}{s s} \frac{\nabla^\mu (\mu/T)}{T} \quad (6)$$

**and should be very small at RHIC because  $\mu/T \sim 0.2$ ,  $n_B/s \sim \mathcal{O}(10^{-3})$**

Consider Bjorken scenario, NO transverse expansion,  $u^\mu(x) = (t, 0, 0, z)/\sqrt{t^2 - z^2}$ , which approximates well the initial evolution in a heavy ion collision, and **follow shear stress only**.

$$T_{LR}^{\mu\nu} = \begin{pmatrix} e & & & \\ & p - \frac{\pi_L}{2} & & \\ & & p - \frac{\pi_L}{2} & \\ & & & p + \pi_L \end{pmatrix} = \begin{pmatrix} e & & & \\ & p_T & & \\ & & p_T & \\ & & & p_L \end{pmatrix}$$

importance of dissipation can be gauged via the pressure anisotropy

$$R \equiv \frac{p_L}{p_T} = \frac{p + \pi_L}{p - \pi_L/2} \quad (\text{typically } \pi_L < 0 \Rightarrow R < 1)$$

**study  $R$  as a function of the initial inverse mean free path  $K_0 \equiv \tau_0/\lambda_{tr,0}$**

**if Grad's approximation is valid, IS should apply at large enough  $K_0$**

take simplest of all cases - 1D Bjorken, massless  $e = 3p$  EOS,  $2 \rightarrow 2$

$$\pi_{LR}^{\mu\nu} = \text{diag}(0, -\frac{\pi_L}{2}, -\frac{\pi_L}{2}, \pi_L), \quad \Pi \equiv 0, \quad q^\mu \equiv 0 \quad (\text{reflection symmetry})$$

$$\dot{p} + \frac{4p}{3\tau} = -\frac{\pi_L}{3\tau} \quad (7)$$

$$\dot{\pi}_L + \frac{\pi_L}{\tau} \left( \frac{2K(\tau)}{3C} + \frac{4}{3} + \frac{\pi_L}{3p} \right) = -\frac{8p}{9\tau}, \quad (8)$$

where


$$K(\tau) \equiv \frac{\tau}{\lambda_{tr}(\tau)}, \quad C \approx \frac{4}{5}. \quad (9)$$

**For  $\sigma = \text{const}$ :**  $\lambda_{tr} = 1/n\sigma_{tr} \propto \tau \Rightarrow K = K_0 = \text{const}, \quad \eta/s \sim T\lambda_{tr} \sim \tau^{2/3}$


**for  $\eta/s \approx \text{const}$ :**  $K = K_0(\tau/\tau_0)^{\sim 2/3} \propto \tau^{\sim 2/3}$

**And we kept the COMPLETE Israel-Stewart equations (every term)**

$\sigma = \text{const}$




$\eta/s \approx \text{const}$



**Need  $K_0 \gtrsim 2 - 3$  for IS hydro to apply, i.e.,  $\lambda_{tr} \lesssim 0.3 - 0.5 \tau_0$**

**In very center of Au+Au at RHIC:  $K_0 \approx 10 - 20$  if  $\eta/s = 1/(4\pi)$**

Same conclusion even if we start from a **LARGE** initial anisotropy  $R \approx 0.3$ , well outside the Navier-Stokes regime.



# Viscous IS hydro in 2D

We solve the full Israel-Stewart equations, including vorticity terms from kinetic theory, in a 2+1D boost-invariant scenario. Shear stress only.

$$\dot{\pi}^{\mu\nu} + \frac{1}{\tau_\pi} \pi^{\mu\nu} = \frac{1}{\beta_2} \nabla^{\langle\mu} u^{\nu\rangle} - \frac{1}{2} \pi^{\mu\nu} D_\alpha u^\alpha - \frac{1}{2} \pi^{\mu\nu} \left[ \ln \frac{\beta_2}{T} \right] + 2 \pi_\lambda^{\langle\mu} \omega^{\nu\rangle\lambda} - (u^\mu \pi^{\nu\alpha} + u^\nu \pi^{\alpha\mu}) \dot{u}_\alpha$$

Mimic a known reliable transport model:

- massless Boltzmann particles  $\Rightarrow \epsilon = 3P$
- only  $2 \leftrightarrow 2$  **processes**, i.e. conserved particle number
- $\eta = 4T/(5\sigma_{tr})$ ,  $\beta_2 = 3/(4p)$
- either  $\sigma = \text{const.} = 47 \text{ mb}$  ( $\sigma_{tr} = 14 \text{ mb}$ )  $\leftarrow$  the simplest in transport  
or  $\sigma \propto \tau^{2/3} \Rightarrow \eta/s \approx 1/(4\pi)$


“RHIC-like” initialization:

- $\tau_0 = 0.6 \text{ fm}/c$
- $b = 8 \text{ fm}$
- $T_0 = 385 \text{ MeV}$  and  $dN/d\eta|_{b=0} = 1000$
- freeze-out at **constant**  $n = 0.365 \text{ fm}^{-3}$

# Pressure evolution in the core

$T^{xx}$  and  $T^{zz}$  averaged over **the core of the system,  $r_{\perp} < 1$  fm:**

$$\eta/s \approx 1/(4\pi) \quad (\sigma \propto \tau^{2/3})$$




Huovinen & DM, arXiv:0806.1367

**remarkable similarity!**

# Viscous hydro elliptic flow

- TWO effects:**
- dissipative corrections to hydro fields  $u^\mu, T, n$
  - dissipative corrections in Cooper-Frye freezeout  $f \rightarrow f_0 + \delta f$

$$\eta/s \approx 1/(4\pi) \quad (\sigma \propto \tau^{2/3})$$



Must use Grad's quadratic correction in Cooper-Frye formula

$$E \frac{dN}{d^3p} = \int p^\mu d\sigma_\mu (f_0 + \delta f)$$


for massless  $\varepsilon = 3p$ , shear only

$$\delta f = f_0 \left[ 1 + \frac{p^\mu p^\nu \pi_{\mu\nu}}{8nT^3} \right]$$

calculation for  $\sigma_{tr} = const \sim 15mb$  shows similar behavior

# Viscous hydro vs transport $v_2$


Huovinen & DM, arXiv:0806.1367




- excellent agreement when  $\sigma = \text{const} \sim 47mb$
- good agreement for  $\eta/s \approx 1/(4\pi)$ , i.e.,  $\sigma \propto \tau^{2/3}$

This means that now all groups agree that viscous corrections to elliptic flow in Au+Au at RHIC are modest  $\sim 20\%$  if  $\eta/s \sim 1/(4\pi)$


Dusling & Teaney, PRC77




Romatschke & Romatschke, arxiv:0706.1522



Huovinen & DM, arXiv:0806.1367






$\sigma \approx 45$  mb result for RHIC corresponds to

$$\eta/s \sim \lambda_{tr} T \sim 1/(\sigma T^2)$$

$$T \sim \tau^{-1/3} \text{ cooling}$$



at early times, violates conjectured viscosity bound DM, arXiv:0806.0026

# Yet more hydro terms?

If we do not start from Israel-Stewart procedure but instead impose conformal invariance (implies  $\varepsilon = 3p$ ), even further terms are possible in the shear stress equation Baier, Romatschke, Son, JHEP04, 100 ('08)

$$\dot{\pi}^{\mu\nu} = \dots + \frac{\lambda_1}{\eta^2} \pi^{\alpha\langle\mu} \pi_{\alpha}^{\nu\rangle} + \lambda_3 \omega^{\alpha\langle\mu} \omega_{\alpha}^{\nu\rangle} \quad (10)$$

In the calculations shown so far,  $\omega$  is rather small, while  $\pi$  should not be very large for hydro to apply. Nevertheless, the importance of these terms depends on the magnitude of matter coefficients in front.

Based on the recent successful hydro-transport comparisons, which did not include the new terms in the hydro, these extra terms are expected to have negligible influence. They matter more, however, for a nonequilibrium theory other than covariant transport.



Note, if we relax conformal invariance, the numerous other terms become possible.

# Bulk viscosity

Recent 0+1D explorations Fries et al, arxiv:0807.4333 based on relaxation eqn

$$\dot{\Pi} = -\frac{1}{\tau_{\Pi}}(\Pi - \Pi_{NS})$$

find significant entropy production:  $\zeta/s_{max} \sim 0.4$  similar to  $\eta/s = 1/4\pi$



# Dissipative hydro - summary

- dissipative hydro describes the evolution of a system **near local equilibrium**, in terms of a few more macroscopic parameters
- causality requires abandoning Navier-Stokes, in favor of second-order formulation, such as Israel-Stewart. This can be motivated both from thermodynamic principles, and from Grad's 14-moment approximation in kinetic theory.
- recent comparison between IS hydro and covariant transport in 0+1D and 2+1D Bjorken geometry shows that the Israel-Stewart (i.e., Grad's 14-moment) approximation, though uncontrolled, is quite accurate in practice
- lot more work ahead - e.g., **latest lattice EOS**, coupling to hadron transport
- **difficult to determine transport coefficients and relaxation times from first principles**
- **conceptual problems with freezeout remain**
- **weakest link, as always, initial conditions** - thermalization mechanism needs to be understood