

# Coordinates, phase space, constraints

Sourendu Gupta

TIFR, Mumbai, India

Classical Mechanics 2011

August 1, 2011

# Mechanics of a single particle

For the motion of a particle (of constant mass  $m$  and position specified by the vector  $\mathbf{x}$ ) viewed from an **inertial frame**, one has the **equations of motion**:

$$m\ddot{\mathbf{x}} = \mathbf{f}, \quad \text{ie,} \quad \dot{\mathbf{x}} = \frac{\mathbf{p}}{m}, \quad \dot{\mathbf{p}} = \mathbf{f}.$$

Here  $\mathbf{f}$  is the **force** acting on the particle and  $\mathbf{p} = m\dot{\mathbf{x}}$  is its **momentum**. The function  $\mathbf{x}(t)$  is the **trajectory** of the particle. This is found with enough **initial conditions**:  $\mathbf{x}(0)$  and  $\mathbf{p}(0)$  are both needed.

## Phase space

The 6-d space with coordinates  $\{\mathbf{x}, \mathbf{p}\}$  is called the **phase space** of the particle. The solutions of the EoM give  $\mathbf{x}(t)$  and  $\mathbf{p}(t)$  which together define a **phase space trajectory**.

# The free particle

## Problem 1

Assume that a particle is constrained to travel along a straight line.

- 1 For such a particle solve the equations of motion when  $f = 0$ .
- 2 What are the trajectories? The phase space trajectories?
- 3 What are the dimensions of **phase space volume**?
- 4 If different particles (with the same equations of motion) have initial conditions all of which are within a small square of phase space at the initial time  $t = 0$ , do the phase space points for the particles all remain within an identical (perhaps shifted) square at a later time?
- 5 Does the (minimum) volume of phase space within which you can find all the above particles change with time?

# The square well potential

## Problem 2

Assume that a particle is constrained to travel along a straight line subject to the potential

$$V(x) = \begin{cases} -V_0 & (|x| \leq a) \\ 0 & (|x| > a) \end{cases}$$

- 1 Solve the equations of motion. What are the trajectories? The phase space trajectories?
- 2 If different particles (with the same equations of motion) have initial conditions all of which are within a small square of phase space at the initial time  $t = 0$ , what happens to this under time evolution?

## A little mathematical generalization

An **ordinary differential equation** of order  $N$  for any vector  $\mathbf{x}_1$

$$\frac{d^N \mathbf{x}_1}{dt^N} = \mathbf{f} \left( t, \mathbf{x}_1, \frac{d\mathbf{x}_1}{dt}, \dots, \frac{d^{N-1} \mathbf{x}_1}{dt^{N-1}} \right),$$

is equivalent to the system of equations

$$\dot{\mathbf{x}}_1 = \mathbf{x}_2, \quad \dot{\mathbf{x}}_2 = \mathbf{x}_3, \quad \dot{\mathbf{x}}_N = \mathbf{f}(t, \mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_{N-1}).$$

One can define a “phase space” with coordinates  $\{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N\}$ . When these are given as functions of time one has a “phase space trajectory”. Clearly this requires complete specification of  $N$  initial conditions, one for each of the  $\mathbf{x}_i$ .

If the force is not an explicit function of time, then the equation is **autonomous**. In an autonomous equation (or system of equations) the phase space trajectories do not intersect. Prove this

# The principle of relativity

## Galileo's principle of relativity

There are frames where the laws of physics are terribly simple: if no forces act on a particle then it remains at rest. This is one inertial frame. Any frame which moves at a constant velocity with respect to this is also an inertial frame.

Here is the transformation law from one inertial frame to another:


$$\mathbf{p}' = \mathbf{p} + m\mathbf{V}, \quad \mathbf{x}' = \mathbf{x} + \mathbf{V}t.$$

## Problem 3


Is a frame fixed with respect to the earth an inertial frame? If not then how badly non-inertial is it?

# Kinds of constraints

- **Non-holonomic constraint.** Example: particle confined in a box. The coordinates are constrained to be  $|\mathbf{x}| \leq L$  where  $L$  is the size of the box. Typically non-holonomic constraints are expressed as inequalities on the coordinates.
- **Holonomic constraint.** Example: particle moving in a line. The coordinates are constrained to be  $\mathbf{x} = s\hat{\mathbf{n}}$  where  $\hat{\mathbf{n}}$  is a unit vector in some direction. Another way to write this is  $\mathbf{x} \times \hat{\mathbf{n}} = 0$ , i.e., the components of  $\mathbf{x}$  orthogonal to  $\hat{\mathbf{n}}$  vanish. Typically holonomic constraints are expressed as equalities.




nonholonomic



holonomic

# Forces of constraint



- Consider the motion of a simple pendulum: a bob suspended from a fixed support by means of an inelastic thread. If the length of the thread is  $L$ , then (if the initial condition is  $\mathbf{p} = 0$ ) the CM of the bob moves along the arc of a circle of radius  $L$ . The thread is in tension. This force of constraint makes the motion one dimensional. There is no work done by the force of constraint. **Prove this**
- In actual fact any thread is elastic, so the length of the pendulum can vary. Similarly, the support is never exactly fixed. In both cases the motion is no longer one-dimensional, and the constraint can do work. A holonomic constraint is a mathematical idealization. **When is it a good idealization?**



# When can a body be considered rigid?

## Problem 4: How rigid is a rigid body?

- ➊ A simple pendulum is hung from a spring with spring constant  $K$ . The motion of the pendulum is a sum of two harmonic motions (one is the swinging of the bob, the other the extension of the spring) assumed not to be in resonance. How does energy flow between these two oscillators? Does the energy flow modify the frequency of the pendulum?
- ➋ A particle tries to bounce off an inflated balloon. Are there any conditions under which there is no bounce? When there is a bounce can the kinetic energy of the particle remain unchanged?
- ➌ Under what conditions does a stone bounce off the surface of water?
- ➍ Is it possible to run on loose sand without slowing down?

## Forces of constraint do no work

A particle is restricted to a surface by a **constraint force**  $\mathbf{R}$ .

**Why not a curve?** The equation of motion gives

$$0 = \mathbf{v} \cdot (m\dot{\mathbf{v}} - \mathbf{R}) = \dot{T} - \mathbf{v} \cdot \mathbf{R},$$

where the **kinetic energy**  $T = mv^2/2$ . If the force of the constraint does no work then  $T$  is conserved and  $\mathbf{v} \cdot \mathbf{R} = 0$ . If the normal to the surface at the point  $\mathbf{x}$  is in the direction  $\hat{\mathbf{n}}$ , then  $\hat{\mathbf{n}} \cdot \mathbf{v} = 0$ , which means that  $\hat{\mathbf{n}} \times \mathbf{R} = 0$ .

If we try to write the constraint force in terms of a potential

$$V' = \sum_{i=1}^3 \alpha_i x_i^2,$$

since the gradient of the potential is zero along the surface, the only non-vanishing component is  $\hat{\mathbf{n}} \cdot \nabla V'$ . The constraint is recovered by taking  $\alpha_i \rightarrow \infty$  subject to this condition.

# Keywords and References

## Keywords

equations of motion, force, inertial frame, kinetic energy, momentum, trajectory, phase space, phase space trajectory, phase space volume, ordinary differential equation, autonomous system of equations, initial conditions, holonomic constraint, non-holonomic constraint, constraint force

## References

Goldstein, chapter 1.

Landau, section 1 and 3.