Re-Creating the Big Bang

Rajiv V. Gavai
T. I. F. R., Mumbai

Re-Creating the Big Bang

Rajiv V. Gavai
T. I. F. R., Mumbai

What is Big Bang ?

Why Re-Create it ?

Heavy Ion Collisions.

Introduction : What is Big Bang ?

Introduction : What is Big Bang ?

- Hubble's Expansion Law. $\Longrightarrow$
  Past : Universe was Denser; Now :
  It is Rarer

Introduction : What is Big Bang ?

- Hubble's Expansion Law. $\Longrightarrow$
  Past : Universe was Denser; Now :
  It is Rarer

- Born in Hot Big Bang; Cooled by Expansion

Introduction : What is Big Bang ?

- Hubble's Expansion Law. $\Longrightarrow$
  Past : Universe was Denser; Now : It is Rarer

- Born in Hot Big Bang; Cooled by Expansion

- Cosmic Microwave Background Radiation (CMBR) — Strongest Evidence.
  - Most perfect black body radiation spectrum.
  - $T \sim 3000^\circ$ K, redshifted due to expansion $T \sim 2.726^\circ$ K.

Figure 1: Our Universe at the age of 380,000 years.

Why Re-Create it ?

Copyright © 2004 Pearson Education, publishing as Addison Wesley.

Strong Interactions

- Known Interactions and Particles a century ago: Electromagnetism, Gravity and Electrons, Atoms.

Strong Interactions

- Known Interactions and Particles a century ago: Electromagnetism, Gravity and Electrons, Atoms.

- Rutherford's Scattering Experiment $\longrightarrow$ various layers that have since been discovered.

Strong Interactions

- Known Interactions and Particles a century ago: Electromagnetism, Gravity and Electrons, Atoms.

- Rutherford's Scattering Experiment $\longrightarrow$ various layers that have since been discovered.

- Quarks and Leptons – Basic building blocks : Proton (uud), Neutron (udd), Pion $(u\bar{d})$....

- A Variety of Vector Bosons : Carriers of forces.

	Gravity	Weak (Electroweak)	Electromagnetic	Strong
Carried By	Graviton (not yet observed)	W^+ W^- Z^0	Photon	Gluon
Acts on	All	Quarks and Leptons	Quarks and Charged Leptons and W^+ W^-	Quarks and Gluons

Strengths in a ratio $10^{-39} : 10^{-5} : 10^{-2} : 1$

	Gravity	Weak (Electroweak)	Electromagnetic	Strong
Carried By	Graviton (not yet observed)	W^+ W^- Z^0	Photon	Gluon
Acts on	All	Quarks and Leptons	Quarks and Charged Leptons and W^+ W^-	Quarks and Gluons

Strengths in a ratio $10^{-39} : 10^{-5} : 10^{-2} : 1$

(Anti-)Quarks come in three (anti-)colours, making gluons also coloured.

Phase Diagram of Strong Matter

- Quantum Chromo Dynamics (QCD) is the (Gauge) Theory of interactions of quarks-gluons.

Phase Diagram of Strong Matter

- Quantum Chromo Dynamics (QCD) is the (Gauge) Theory of interactions of quarks-gluons.
- Much richer structure : Quark Confinement, Dynamical Symmetry Breaking..
- New States at High Temperatures/Density expected on basis of models.

Phase Diagram of Strong Matter

- Quantum Chromo Dynamics (QCD) is the (Gauge) Theory of interactions of quarks-gluons.
- Much richer structure : Quark Confinement, Dynamical Symmetry Breaking..
- New States at High Temperatures/Density expected on basis of models.
- Quark-Gluon Plasma, such a new phase, expected in Relativistic Heavy Ion Collisions.
- Color Superconductivity another phase, may exist in very dense stars.

<L> × FUKUGITA + UKAWA
<ψ̄ψ> □ FUKUGITA + UKAWA
LANGEVIN
<L> ▽ POLONYI ET AL.
<ψ̄ψ> ○ POLONYI ET AL.
MICROCANONICAL
<L> ▼ GAVAI
<ψ̄ψ> ● GAVAI
PSEUDO-FERMION
<L>
<ψ̄ψ>
8³ x 4 LATTICE
ma = 0.1
4 FLAVOURS
1.2
1.0
0.8
0.6
0.4
0.2
3.5
4.5
5.5
6.5
Tc
β = 6/g²

Temperature T [MeV]
200
100
0
Early universe
Quarks and Gluons
Deconfinement & chiral transition
Critical point?
RHIC & LHC
Hadrons
GSI SIS 200
Color Super-conductor?
Neutron stars
1
Nuclei
Net Baryon Density

- The Transition Temperature $T_c \sim 175$ MeV (about 2 Trillion $^\circ$ K).

- T_c, the Equation of State (EOS) and many other properties, notably the Wróblewski Parameter λ_s and other correlations for Heavy Ion Physics have been predicted theoretically.

Gavai and Gupta, Phys Rev D65, 2002 and Phys.Rev. D73, 2006.

Gavai and Gupta, Phys Rev D65, 2002 and Phys.Rev. D73, 2006.

- λ_s — Measure of Production of strange quark-antiquark pairs; Expts agree with estimates from the new state Quark-Gluon Plasma.
  — Lattice QCD suggests that strangeness carried by quark-like objects
  — Generally flavour shows quasi-quark behaviour.

Our Workhorse

CRAY X1 of I L G T I , T I F R, Mumbai

Our Results

1.1
1
0.9
0.8
0.7
T/Tc
0
1
2
3
4
5
μB/T
18 GeV (CERN)
30 GeV
20 GeV
10 GeV
Freezeout curve

Our Results

- We (RVG & S. Gupta, PRD 2005) find the Critical Point at smaller $\mu_B/T \sim 1$.
- Our estimate consistent with Fodor & Katz (2002) [$m_\pi/m_\rho = 0.31$ and $N_s m_\pi \sim$ 3-4].

Our Results

- We (RVG & S. Gupta, PRD 2005) find the Critical Point at smaller $\mu_B/T \sim 1$.
- Our estimate consistent with Fodor & Katz (2002) [$m_\pi/m_\rho = 0.31$ and $N_s m_\pi \sim$ 3-4].
- Strong finite size effects for small N_s. A strong change around $N_s m_\pi \sim 6$.
- RHIC, if run at lower energy, can potentially discover it.

Heavy Ion Collisions

- Where does one find these new phases ? Can they be produced in laboratory ?

Heavy Ion Collisions

- Where does one find these new phases ? Can they be produced in laboratory ?
- Early Universe — About $10 - 20\mu$s after the Big Bang and in Cores of Dense Neutron Stars

Heavy Ion Collisions

- Where does one find these new phases ? Can they be produced in laboratory ?
- Early Universe — About $10 - 20\mu$s after the Big Bang and in Cores of Dense Neutron Stars
- Quark-Gluon Plasma can be, and may **indeed** have been, produced in Heavy Ion Collisions in CERN, Geneva and BNL, New York.

Heavy Ion Collisions

- Where does one find these new phases ? Can they be produced in laboratory ?
- Early Universe — About $10 - 20\mu$s after the Big Bang and in Cores of Dense Neutron Stars
- Quark-Gluon Plasma can be, and may **indeed** have been, produced in Heavy Ion Collisions in CERN, Geneva and BNL, New York.
- Necessary Conditions for QGP production :
  - High Energy Density, $\approx$ 1-3 GeV/fm^3.
  - Large System Size, $L \gg \Lambda_{QCD}^{-1}$.
  - Many particles

$\Longrightarrow$ Heavy Ion Collisions at 99.5-99.995 % Velocity of Light.

$\Longrightarrow$ Heavy Ion Collisions at 99.5-99.995 % Velocity of Light.

$\Longrightarrow$ Heavy Ion Collisions at 99.5-99.995 % Velocity of Light.

$\Longrightarrow$ Heavy Ion Collisions at 99.5-99.995 % Velocity of Light.

Fireball of QGP condenses into hadrons in $\approx 10^{-23}$ seconds.

How to look for QGP

- Jet Quenching :

How to look for QGP

- Jet Quenching :
  - Rare, Highly Energetic Scatterings produce jets of particles : $g + g \rightarrow g + g$.
  - Quark-Gluon Plasma, any medium in general, interacts with a jet, causing it to lose energy – Jet Quenching.

How to look for QGP

- Jet Quenching :
  - Rare, Highly Energetic Scatterings produce jets of particles : $g + g \rightarrow g + g$.
  - Quark-Gluon Plasma, any medium in general, interacts with a jet, causing it to lose energy – Jet Quenching.
  - On-Off test possible – Compare Collisions of Heavy-Heavy nuclei with Light-Heavy or Light-Light.

Azimuthal distributions

Near- side: p+p, d+Au, Au+Au similar

Back- to- back: Au+Au strongly suppressed relative to p+p and d+Au

Azimuthal distributions

Near- side: p+p, d+Au, Au+Au similar

Back- to- back: Au+Au strongly suppressed relative to p+p and d+Au

- Anisotropic Flow & QGP as Perfect Liquid
- Debye Screening of Quarks $\Longrightarrow$ No binding to Hadrons — Anomalous J/ψ Suppression

Summary

- Lattice QCD **predicts** new states of strongly interacting matter and is able to shed light on the properties of the Quark-Gluon plasma phase.

- Our results on correlations of quantum numbers suggest QGP to have quarklike excitations.

Summary

- Lattice QCD **predicts** new states of strongly interacting matter and is able to shed light on the properties of the Quark-Gluon plasma phase.

- Our results on correlations of quantum numbers suggest QGP to have quarklike excitations.

- Phase diagram in $T - \mu_B$ plane has begun to emerge: Our estimate for the critical point is $\mu_B/T \sim 1 - 2$.

Summary

- Lattice QCD **predicts** new states of strongly interacting matter and is able to shed light on the properties of the Quark-Gluon plasma phase.

- Our results on correlations of quantum numbers suggest QGP to have quarklike excitations.

- Phase diagram in $T - \mu_B$ plane has begun to emerge: Our estimate for the critical point is $\mu_B/T \sim 1 - 2$.

Heavy Ion Collisions in CERN Geneva, and BNL, New York, have seen tell-tale signs of QGP : Many surprises already and more excitement likely to come.